

Datasheet for ABIN7563871  
**IKK alpha Protein (AA 1-745) (His tag)**



[Go to Product page](#)

## Overview

Quantity:	1 mg
Target:	IKK alpha (CHUK)
Protein Characteristics:	AA 1-745
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IKK alpha protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

## Product Details

Purpose:	Custom-made recombinat Chuk Protein expressed in mammalien cells.
Sequence:	MERPPGLRPG AGGPWEMRER LGTGGFGNVS LYQHRELDLK IAIKSCRLEL SSKNRERWCH EIQIMKKLDH ANVVKACDVP EELNFLINDV PLLAMEYCSG GDLRKLINPK ENCCGLKESQ ILSLLSDIGS GIRYLHENKI IHRDLKPENI VLQDVGGKTI HKIIDLGYAK DVDQGSLLCTS FVGTLQYLAP ELFENKPYTA TVDYWSFGTM VFECIAGYRP FLHHLQPFTW HEKIKKKDPK CIFACEEMTG EVRFSSHLPQ PNSLCSLIVE PMESWLQLML NWDPPQRRGGP IDLTLKQPRC FALMDHILNL KIVHILNMTS AKIISFLLPC DESLHSLQSR IERETGINTG SQELLSETGI SLDPRKPASQ CVLDGVRGCD SYMVYLFDKS KTVYEGPFAS RSLSDCVNYI VQDSKIQLPI IQLRKVWAEA VHYVSGLKED YSRLFQGQRA AMLSLLRYNA NLTKMKNTLI SASQQLKAKL EFFRKSQILD LERYSEQMTY GISSEKMLKA WKEMEEKAIH YSEVGVIGYL EDQIMSLHTE IMELQKSPYG RRQGDLMESL EQRAIDLYKQ LKHRPPDHLY SDSTEMVKII VHTVQSQDRV LKELFGHLSK LLGCKQKIID LLPKVEVALS NIKEADNTVM FMQGRQKEI WHLLKIACTQ

## Product Details

---

SSARSLVGSS LEGTVTPPVV AWLPPTLADR EHPLTCVVTP QDGETLAQMI EENLNCLGHL  
STIIREANED QSSSLMSLDW SWLAE **Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

---

### Characteristics:

#### Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

---

### Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

---

### Grade:

custom-made

---

## Target Details

---

### Target:

IKK alpha (CHUK)

---

### Alternative Name:

Chuk ([CHUK Products](#))

---

### Background:

Inhibitor of nuclear factor kappa-B kinase subunit alpha (I-kappa-B kinase alpha) (IKK-A) (IKK-alpha) (Ikbka) (IkkappaB kinase) (EC 2.7.11.10) (Conserved helix-loop-helix ubiquitous kinase) (I-kappa-B kinase 1) (IKK1) (Nuclear factor NF-kappa-B inhibitor kinase alpha) (NFKB1A),FUNCTION: Serine kinase that plays an essential role in the NF-kappa-B signaling pathway which is activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA damages or other cellular stresses. Acts as a part of the canonical IKK complex in the conventional pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B on serine residues. These modifications allow polyubiquitination of the inhibitors and subsequent degradation by the proteasome. In turn, free NF-kappa-B is translocated into the

## Target Details

---

nucleus and activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis. Negatively regulates the pathway by phosphorylating the scaffold protein TAXBP1 and thus promoting the assembly of the A20/TNFAIP3 ubiquitin-editing complex (composed of A20/TNFAIP3, TAX1BP1, and the E3 ligases ITCH and RNF11). Therefore, CHUK plays a key role in the negative feedback of NF-kappa-B canonical signaling to limit inflammatory gene activation. As part of the non-canonical pathway of NF-kappa-B activation, the MAP3K14-activated CHUK/IKKA homodimer phosphorylates NFKB2/p100 associated with RelB, inducing its proteolytic processing to NFKB2/p52 and the formation of NF-kappa-B RelB-p52 complexes. In turn, these complexes regulate genes encoding molecules involved in B-cell survival and lymphoid organogenesis. Participates also in the negative feedback of the non-canonical NF-kappa-B signaling pathway by phosphorylating and destabilizing MAP3K14/NIK. Within the nucleus, phosphorylates CREBBP and consequently increases both its transcriptional and histone acetyltransferase activities. Modulates chromatin accessibility at NF-kappa-B-responsive promoters by phosphorylating histones H3 at 'Ser-10' that are subsequently acetylated at 'Lys-14' by CREBBP (PubMed:12789342). Additionally, phosphorylates the CREBBP-interacting protein NCOA3. Also phosphorylates FOXO3 and may regulate this pro-apoptotic transcription factor. Phosphorylates RIPK1 at 'Ser-25' which represses its kinase activity and consequently prevents TNF-mediated RIPK1-dependent cell death (PubMed:30988283). Phosphorylates AMBRA1 following mitophagy induction, promoting AMBRA1 interaction with ATG8 family proteins and its mitophagic activity (By similarity). {ECO:0000250|UniProtKB:O15111, ECO:0000269|PubMed:12789342, ECO:0000269|PubMed:30988283}.

---

Molecular Weight: 84.7 kDa

---

UniProt: [Q60680](#)

---

Pathways: [PI3K-Akt Signaling](#), [NF-kappaB Signaling](#), [RTK Signaling](#), [TCR Signaling](#), [TLR Signaling](#), [Fc-epsilon Receptor Signaling Pathway](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#), [Activation of Innate immune Response](#), [Hepatitis C](#), [Toll-Like Receptors Cascades](#), [BCR Signaling](#), [Ubiquitin Proteasome Pathway](#), [S100 Proteins](#)

## Application Details

---

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

---

Restrictions: For Research Use only

## Handling

---

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months