

Datasheet for ABIN7563878
TRAF3 Protein (AA 1-567) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	TRAF3
Protein Characteristics:	AA 1-567
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TRAF3 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Purpose:	Custom-made recombinat Traf3 Protein expressed in mammalian cells.
Sequence:	<p>MESSKKMDAA GTLQPNPPLK LQPDRGAGSV LVPEQGGYKE KFKVTVEDKY KCEKRLVLC</p> <p>NPKQTECGHR FCESCMAALL SSSSPKCTAC QESIICKDVF KDNCKREIL ALQVYCRNEG</p> <p>RGCAEQLTLG HLLVHLKNEC QFEELPCLRA DCKEVLKRD LRDHVEKACK YREATCSHCK</p> <p>SQVPMIKLQK HEDTDCPCVV VSCPHKCSVQ TLLRSELSAH LSECVNAPST CSFKRYGCVF</p> <p>QGTNQQIKAH EASSAVQHVN LLKEWSNSLE KKVSLQNES VEKNKSIQSL HNQICSFEIE</p> <p>IERQKEMLRN NESKILHLQR VIDSQAEKLE ELDKEIRPFR QNWEEADSMK SSVESLQNRV</p> <p>TELESVDKSA GQAARNTGLL ESQLSRHDQM LSVHDIRLAD MDLRFQVLET ASYNGVLIWK</p> <p>IRDYKRRKQE AVMGKTLSTLY SQPFYTYGYFG YKMCARVYLN GDGMGKGTHL SLFFVIMRGE</p> <p>YDALLPWPFK QKVTLMLMDQ GSSRRHLGDA FKPDPNSSSF KKPTGEMNIA SGCPVFVAQT</p> <p>VLENGTYIKD DTIFIKVIVD TSDLDPD Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein</p>

could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target:

TRAF3

Alternative Name:

Traf3 ([TRAF3 Products](#))

Background:

TNF receptor-associated factor 3 (EC 2.3.2.27) (CD40 receptor-associated factor 1) (CRAF1) (CD40-binding protein) (CD40BP) (RING-type E3 ubiquitin transferase TRAF3), FUNCTION: Cytoplasmic E3 ubiquitin ligase that regulates various signaling pathways, such as the NF-kappa-B, mitogen-activated protein kinase (MAPK) and interferon regulatory factor (IRF) pathways, and thus controls a lot of biological processes in both immune and non-immune cell types (PubMed:17015635). In TLR and RLR signaling pathways, acts as an E3 ubiquitin ligase promoting the synthesis of 'Lys-63'-linked polyubiquitin chains on several substrates such as ASC that lead to the activation of the type I interferon response or the inflammasome (PubMed:19898473, PubMed:23871208, PubMed:26305951, PubMed:23150880). Following the activation of certain TLRs such as TLR4, acts as a negative NF-kappa-B regulator, possibly to avoid unregulated inflammatory response, and its degradation via 'Lys-48'-linked polyubiquitination is required for MAPK activation and production of inflammatory cytokines

Target Details

(PubMed:16306937). Alternatively, when TLR4 orchestrates bacterial expulsion, TRAF3 undergoes 'Lys-33'-linked polyubiquitination and subsequently binds to RALGDS, mobilizing the exocyst complex to rapidly expel intracellular bacteria back for clearance. Acts also as a constitutive negative regulator of the alternative NF-kappa-B pathway, which controls B-cell survival and lymphoid organ development (PubMed:17723217). Required for normal antibody isotype switching from IgM to IgG (PubMed:19228877). Plays a role T-cell dependent immune responses (PubMed:8934568). Down-regulates proteolytic processing of NFKB2, and thereby inhibits non-canonical activation of NF-kappa-B. Promotes ubiquitination and proteasomal degradation of MAP3K14. {ECO:0000269|PubMed:16306936, ECO:0000269|PubMed:16306937, ECO:0000269|PubMed:17015635, ECO:0000269|PubMed:17158868, ECO:0000269|PubMed:17723217, ECO:0000269|PubMed:18313334, ECO:0000269|PubMed:18997792, ECO:0000269|PubMed:18997794, ECO:0000269|PubMed:19228877, ECO:0000269|PubMed:19898473, ECO:0000269|PubMed:26305951, ECO:0000269|PubMed:26474655, ECO:0000269|PubMed:30579117, ECO:0000269|PubMed:8934568}.

Molecular Weight: 64.3 kDa

UniProt: [Q60803](#)

Pathways: [NF-kappaB Signaling](#), [Apoptosis](#), [TLR Signaling](#), [Activation of Innate immune Response](#), [Hepatitis C](#), [Toll-Like Receptors Cascades](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Handling

Expiry Date: 12 months