

## Datasheet for ABIN7564329

# B3GALNT2 Protein (AA 1-504) (His tag)



### Overview

Quantity:	1 mg
Target:	B3GALNT2
Protein Characteristics:	AA 1-504
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This B3GALNT2 protein is labelled with His tag.

Product Details	
Purpose:	Custom-made recombinant B3gaInt2 Protein expressed in mammalian cells.
Sequence:	MRNWLVLLCP CVLGAALHLW HLWLRSPPDP HNTGPSAADQ SALFPHWKFS HYDVVVGVLS
	ARNNHELRNV IRNTWLKNLL HHPTLSQRVL VKFIIGARGC EVPVEDREDP YSCRLLNITN
	PVLNQEIEAF SFPEDASSSR LSEDRVVSVS FRVLYPIVIT SLGVFYDASD VGFQRNITVK
	LYQTEQEEAL FIARFSPPSC GVQVNKLWYK PVEQFILPES FEGTIVWESQ DLHGLVSRNL
	HRVTVNDGGG VLRVLAAGEG ALPHEFMEGV EGVAGGFIYT VQEGDALLRS LYSRPQRLAD
	HIQDLQVEDA LLQEESSVHD DIVFVDVVDT YRNVPAKLLN FYRWTVESTS FDLLLKTDDD
	CYIDLEAVFN RIAQKNLDGP NFWWGNFRLN WAVDRTGKWQ ELEYPSPAYP AFACGSGYVI
	SKDIVDWLAG NSRRLKTYQG EDVSMGIWMA AIGPKRHQDS LWLCEKTCET GMLSSPQYSP
	EELSKLWELK ELCGDPCQCE AKVR Sequence without tag. The proposed Purification-Tag is
	based on experiences with the expression system, a different complexity of the protein
	could make another tag necessary. In case you have a special request, please contact us.
Specificity:	If you are looking for a specific domain and are interested in a partial protein or a different

#### **Product Details**

	isoform, please contact us regarding an individual offer.
Characteristics:	Key Benefits:
	Made to order protein - from design to production - by highly experienced protein experts.
	Protein expressed in mammalian cells and purified in one-step affinity chromatography
	The optimized expression system ensures reliability for intracellular, secreted and
	<ul><li>transmembrane proteins.</li><li>State-of-the-art algorithm used for plasmid design (Gene synthesis).</li></ul>
	State-of-the-art algorithm used for plasmid design (defie synthesis).
	This protein is a made-to-order protein and will be made for the first time for your order. Our
	experts in the lab try to ensure that you receive soluble protein.
	If you are not interested in a full length protein, please contact us for individual protein
	fragments.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom
	made proteins from other companies is that there is no financial obligation in case the protein
	cannot be expressed or purified.
Purity:	> 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)
Grade:	custom-made
Target Details	
Target:	B3GALNT2
Altornative News	
Alternative Name:	B3gaInt2 (B3GALNT2 Products)
	UDP-GalNAc:beta-1,3-N-acetylgalactosaminyltransferase 2 (Beta-1,3-GalNAc-T2) (EC 2.4.1.313)
	UDP-GalNAc:beta-1,3-N-acetylgalactosaminyltransferase 2 (Beta-1,3-GalNAc-T2) (EC 2.4.1.313) (Beta-1,3-N-acetylgalactosaminyltransferase II),FUNCTION: Beta-1,3-N-acetylgalactosaminyltransferase that synthesizes a unique carbohydrate structure, GalNAc-
	UDP-GalNAc:beta-1,3-N-acetylgalactosaminyltransferase 2 (Beta-1,3-GalNAc-T2) (EC 2.4.1.313) (Beta-1,3-N-acetylgalactosaminyltransferase II),FUNCTION: Beta-1,3-N-acetylgalactosaminyltransferase that synthesizes a unique carbohydrate structure, GalNAc-
	UDP-GalNAc:beta-1,3-N-acetylgalactosaminyltransferase 2 (Beta-1,3-GalNAc-T2) (EC 2.4.1.313) (Beta-1,3-N-acetylgalactosaminyltransferase II),FUNCTION: Beta-1,3-N-acetylgalactosaminyltransferase that synthesizes a unique carbohydrate structure, GalNAc-beta-1-3GlcNAc, on N- and O-glycans. Has no galactose nor galactosaminyl transferase activity toward any acceptor substrate. Involved in alpha-dystroglycan (DAG1) glycosylation: acts
	UDP-GalNAc:beta-1,3-N-acetylgalactosaminyltransferase 2 (Beta-1,3-GalNAc-T2) (EC 2.4.1.313) (Beta-1,3-N-acetylgalactosaminyltransferase II),FUNCTION: Beta-1,3-N-acetylgalactosaminyltransferase that synthesizes a unique carbohydrate structure, GalNAc-beta-1-3GlcNAc, on N- and O-glycans. Has no galactose nor galactosaminyl transferase activity toward any acceptor substrate. Involved in alpha-dystroglycan (DAG1) glycosylation: acts
	UDP-GalNAc:beta-1,3-N-acetylgalactosaminyltransferase 2 (Beta-1,3-GalNAc-T2) (EC 2.4.1.313) (Beta-1,3-N-acetylgalactosaminyltransferase II),FUNCTION: Beta-1,3-N-acetylgalactosaminyltransferase that synthesizes a unique carbohydrate structure, GalNAc-beta-1-3GlcNAc, on N- and O-glycans. Has no galactose nor galactosaminyl transferase activity toward any acceptor substrate. Involved in alpha-dystroglycan (DAG1) glycosylation: acts coordinately with GTDC2/POMGnT2 to synthesize a GalNAc-beta3-GlcNAc-beta-terminus at the
Alternative Name:  Background:	UDP-GalNAc:beta-1,3-N-acetylgalactosaminyltransferase 2 (Beta-1,3-GalNAc-T2) (EC 2.4.1.313) (Beta-1,3-N-acetylgalactosaminyltransferase II),FUNCTION: Beta-1,3-N-acetylgalactosaminyltransferase that synthesizes a unique carbohydrate structure, GalNAc-beta-1-3GlcNAc, on N- and O-glycans. Has no galactose nor galactosaminyl transferase activity toward any acceptor substrate. Involved in alpha-dystroglycan (DAG1) glycosylation: acts coordinately with GTDC2/POMGnT2 to synthesize a GalNAc-beta3-GlcNAc-beta-terminus at the 4-position of protein O-mannose in the biosynthesis of the phosphorylated O-mannosyl
	UDP-GalNAc:beta-1,3-N-acetylgalactosaminyltransferase 2 (Beta-1,3-GalNAc-T2) (EC 2.4.1.313) (Beta-1,3-N-acetylgalactosaminyltransferase II),FUNCTION: Beta-1,3-N-acetylgalactosaminyltransferase that synthesizes a unique carbohydrate structure, GalNAc-beta-1-3GlcNAc, on N- and O-glycans. Has no galactose nor galactosaminyl transferase activity toward any acceptor substrate. Involved in alpha-dystroglycan (DAG1) glycosylation: acts coordinately with GTDC2/POMGnT2 to synthesize a GalNAc-beta3-GlcNAc-beta-terminus at the 4-position of protein O-mannose in the biosynthesis of the phosphorylated O-mannosyl trisaccharide (N-acetylgalactosamine-beta-3-N-acetylglucosamine-beta-4-(phosphate-6-
	UDP-GalNAc:beta-1,3-N-acetylgalactosaminyltransferase 2 (Beta-1,3-GalNAc-T2) (EC 2.4.1.313) (Beta-1,3-N-acetylgalactosaminyltransferase II),FUNCTION: Beta-1,3-N-acetylgalactosaminyltransferase that synthesizes a unique carbohydrate structure, GalNAc-beta-1-3GlcNAc, on N- and O-glycans. Has no galactose nor galactosaminyl transferase activity toward any acceptor substrate. Involved in alpha-dystroglycan (DAG1) glycosylation: acts coordinately with GTDC2/POMGnT2 to synthesize a GalNAc-beta3-GlcNAc-beta-terminus at the 4-position of protein O-mannose in the biosynthesis of the phosphorylated O-mannosyl trisaccharide (N-acetylgalactosamine-beta-3-N-acetylglucosamine-beta-4-(phosphate-6-)mannose), a carbohydrate structure present in alpha-dystroglycan, which is required for

## **Target Details** UniProt: Q8BG28 **Application Details** We expect the protein to work for functional studies. As the protein has not been tested for Application Notes: functional studies yet we cannot offer a guarantee though. Restrictions: For Research Use only Handling Format: Liquid Buffer: The buffer composition is at the discretion of the manufacturer. Avoid repeated freeze-thaw cycles. Handling Advice: -80 °C Storage:

Storage Comment:

Expiry Date:

Store at -80°C.

12 months