

Datasheet for ABIN7564348  
**FTO Protein (AA 1-502) (His tag)**



[Go to Product page](#)

## Overview

Quantity:	1 mg
Target:	FTO
Protein Characteristics:	AA 1-502
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This FTO protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

## Product Details

Purpose:	Custom-made recombinat Fto Protein expressed in mammalian cells.
Sequence:	<p>MKRVQTAEER EREAKKLRL EEDTWPY LTPKDDEFYQ QWQLKYPKLV FREAGSIPEE</p> <p>LHKEVPEAFL TLHKHGCLFR DVVRIQGKDV LTPVSRILIG DPGCTYKYN TRLFVWPWPV</p> <p>KGCTVKYTEA EIAAACQTFL KLNDYLQVET IQALEELAVR EKANEDAVPL CMAEFPRAGV</p> <p>GPSCDDEVDL KSRAAYNVTL LNFMDPQKMP YLKEEPYFGM GKMAVSWHHD ENLVDRSAVA</p> <p>VYSYSCGESE DESEDESSFE GRDPDTWHVG FKISWDIETP GLTIPLHQGD CYFMLDDLNA</p> <p>THQHCVLGAS QPRFSSTHRV AECSTGTLDY ILERCQLALQ NVLNSDDGD VSLKSFDPAV</p> <p>LKQGEEIHNE VEFWLRQFW FQGNRYKLCT DWWCEPMTHL EGLWKKMESM TNAVLREVKR</p> <p>EGLPVEQRSE ILSAILVPLT VRQNLKKEWH ARCQSRVVRT LPVQKPKDCR PYWEKDDPSM</p> <p>PLPFDLTDVV SELRGQLLEA RS <b>Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</b></p>

## Product Details

### Characteristics:

### Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

### Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

### Grade:

custom-made

## Target Details

### Target:

FTO

### Alternative Name:

Fto ([FTO Products](#))

### Background:

Alpha-ketoglutarate-dependent dioxygenase FTO (Fat mass and obesity-associated protein) (Protein fatso) (U6 small nuclear RNA (2'-O-methyladenosine-N(6)-)-demethylase FTO) (EC 1.14.11.-) (U6 small nuclear RNA N(6)-methyladenosine-demethylase FTO) (EC 1.14.11.-) (mRNA (2'-O-methyladenosine-N(6)-)-demethylase FTO) (m6A(m)-demethylase FTO) (EC 1.14.11.-) (mRNA N(6)-methyladenosine demethylase FTO) (EC 1.14.11.53) (tRNA N1-methyl adenine demethylase FTO) (EC 1.14.11.-),FUNCTION: RNA demethylase that mediates oxidative demethylation of different RNA species, such as mRNAs, tRNAs and snRNAs, and acts as a regulator of fat mass, adipogenesis and energy homeostasis (PubMed:17991826, PubMed:18775698, PubMed:28002401). Specifically demethylates N(6)-methyladenosine (m6A) RNA, the most prevalent internal modification of messenger RNA (mRNA) in higher eukaryotes (PubMed:28002401). M6A demethylation by FTO affects mRNA expression and stability (By similarity). Also able to demethylate m6A in U6 small nuclear RNA (snRNA) (By similarity). Mediates demethylation of N(6),2'-O-dimethyladenosine cap (m6A(m)), by

## Target Details

demethylating the N(6)-methyladenosine at the second transcribed position of mRNAs and U6 snRNA (PubMed:28002401). Demethylation of m6A(m) in the 5'-cap by FTO affects mRNA stability by promoting susceptibility to decapping (By similarity). Also acts as a tRNA demethylase by removing N(1)-methyladenine from various tRNAs (By similarity). Has no activity towards 1-methylguanine (By similarity). Has no detectable activity towards double-stranded DNA (By similarity). Also able to repair alkylated DNA and RNA by oxidative demethylation: demethylates single-stranded RNA containing 3-methyluracil, single-stranded DNA containing 3-methylthymine and has low demethylase activity towards single-stranded DNA containing 1-methyladenine or 3-methylcytosine (PubMed:17991826, PubMed:18775698). Ability to repair alkylated DNA and RNA is however unsure in vivo (PubMed:17991826, PubMed:18775698). Involved in the regulation of fat mass, adipogenesis and body weight, thereby contributing to the regulation of body size and body fat accumulation (PubMed:19234441, PubMed:19680540, PubMed:21076408, PubMed:23817550, PubMed:23300482). Involved in the regulation of thermogenesis and the control of adipocyte differentiation into brown or white fat cells (PubMed:19234441, PubMed:19680540). Regulates activity of the dopaminergic midbrain circuitry via its ability to demethylate m6A in mRNAs (PubMed:23817550). {ECO:0000250|UniProtKB:Q9C0B1, ECO:0000269|PubMed:17991826, ECO:0000269|PubMed:18775698, ECO:0000269|PubMed:19234441, ECO:0000269|PubMed:19680540, ECO:0000269|PubMed:21076408, ECO:0000269|PubMed:23300482, ECO:0000269|PubMed:23817550, ECO:0000269|PubMed:28002401}.

Molecular Weight:	58.0 kDa
-------------------	----------

UniProt:	<a href="#">Q8BGW1</a>
----------	------------------------

## Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Liquid
---------	--------

Buffer:	The buffer composition is at the discretion of the manufacturer.
---------	--

## Handling

---

Handling Advice:	Avoid repeated freeze-thaw cycles.
------------------	------------------------------------

Storage:	-80 °C
----------	--------

Storage Comment:	Store at -80°C.
------------------	-----------------

Expiry Date:	12 months
--------------	-----------