

Datasheet for ABIN7564659

BRE Protein (AA 1-383) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	BRE
Protein Characteristics:	AA 1-383
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This BRE protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Purpose:	Custom-made recombinant Babam2 Protein expressed in mammalian cells.
Sequence:	<p>MSPEIALNRI SPMLSPFISS VVRNGKVGLD ATNCLRITDL KSGCTSLTPG PNCDFKLHI</p> <p>PYAGETLKWD IIFNAQYPEL PPDFIFGEDA EFLPDPSALH NLASWNPSNP ECLLLVVKEL</p> <p>VQQYHQFQCG RLRESSRLMF EYQTLLEEPQ YGENMEIYAG KKNWWTGEFS ARFLKL PVD</p> <p>FSNIPTYLLK DVNEDPGEDV ALLSVSFEDT EATQVYPKLY LSPRIEHALG GSSALHIPAF</p> <p>PGGGCLIDYV PQVCHLLTNK VQYVIQGYHK RREYIAAFLS HFGTGVVEYD AEGFTKL TLL</p> <p>LMWKDFCFLV HIDLPLFFPR DQPTLTFQSV YHFTNSGQLY SQAQKNYPYS PRWDGNEMAK</p> <p>RAKAYFKTFV PQFQEAFAFAN GKL Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p>
Characteristics:	Key Benefits:

Product Details

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:	> 90 % as determined by Bis-Tris Page, Western Blot
---------	---

Grade:	custom-made
--------	-------------

Target Details

Target:	BRE
---------	-----

Alternative Name:	Babam2 (BRE Products)
-------------------	---

Background:	<p>BRISC and BRCA1-A complex member 2 (BRCA1-A complex subunit BRE) (BRCA1/BRCA2-containing complex subunit 45) (Brain and reproductive organ-expressed protein),FUNCTION: Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it acts as an adapter that bridges the interaction between BABAM1/NBA1 and the rest of the complex, thereby being required for the complex integrity and modulating the E3 ubiquitin ligase activity of the BRCA1-BARD1 heterodimer. Probably also plays a role as a component of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin (By similarity). May regulate TNF-alpha signaling through its interactions with TNFRSF1A. {ECO:0000250, ECO:0000269 PubMed:9737713}, FUNCTION: Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions</p>
-------------	--

Target Details

sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it acts as an adapter that bridges the interaction between BABAM1/NBA1 and the rest of the complex, thereby being required for the complex integrity and modulating the E3 ubiquitin ligase activity of the BRCA1-BARD1 heterodimer. Component of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates. Within the BRISC complex, acts as an adapter that bridges the interaction between BABAM1/NBA1 and the rest of the complex, thereby being required for the complex integrity. The BRISC complex is required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitinating NUMA1. The BRISC complex plays a role in interferon signaling via its role in the deubiquitination of the interferon receptor IFNAR1, deubiquitination increases IFNAR1 activity by enhancing its stability and cell surface expression. Down-regulates the response to bacterial lipopolysaccharide (LPS) via its role in IFNAR1 deubiquitination. May play a role in homeostasis or cellular differentiation in cells of neural, epithelial and germline origins (By similarity). May also act as a death receptor-associated anti-apoptotic protein, which inhibits the mitochondrial apoptotic pathway. May regulate TNF-alpha signaling through its interactions with TNFRSF1A, however these effects may be indirect (PubMed:9737713). {ECO:0000250|UniProtKB:Q9NXR7, ECO:0000305|PubMed:9737713}.

Molecular Weight:	43.5 kDa
UniProt:	Q8K3W0
Pathways:	Positive Regulation of Response to DNA Damage Stimulus

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.

Handling

Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months