

Datasheet for ABIN7564777
EGLN3 Protein (AA 1-239) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	EGLN3
Protein Characteristics:	AA 1-239
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This EGLN3 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Purpose:	Custom-made recombinat EglN3 Protein expressed in mammalien cells.
Sequence:	<p>MPLGHIMRLD LEKIALEYIV PCLHEVGFCY LDNFLGEVVG DCVLERVKQL HYNALRDGQ LAGPRAGVSK RHLRGDQITW IGGNEEGCEA INFLSLIDR LVLYCGSRLG KYVVKERSKA MVACYPGNGT GYVRHVDNPN GDGRCITCIY YLNKNWDAKL HGGVLRIFPE GKSFVADVEP IFDRLLFFWS DRRNPHEVQP SYATRYAMTV WYFDAEERAE AKKKFRNLTR KTESALAKD</p> <p>Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary.</p> <p>In case you have a special request, please contact us.</p>
Characteristics:	<p>Key Benefits:</p> <ul style="list-style-type: none"> • Made to order protein - from design to production - by highly experienced protein experts. • Protein expressed in mammalien cells and purified in one-step affinity chromatography • The optimized expression system ensures reliability for intracellular, secreted and

transmembrane proteins.

- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity: > 90 % as determined by Bis-Tris Page, Western Blot

Grade: custom-made

Target Details

Target: EGLN3

Alternative Name: EglN3 (EGLN3 Products)

Background: Prolyl hydroxylase EGLN3 (EC 1.14.11.-) (Egl nine homolog 3) (EC 1.14.11.29) (Hypoxia-inducible factor prolyl hydroxylase 3) (HIF-PH3) (HIF-prolyl hydroxylase 3) (HPH-3) (Prolyl hydroxylase domain-containing protein 3) (PHD3) (SM-20),FUNCTION: Prolyl hydroxylase that mediates hydroxylation of proline residues in target proteins, such as PKM, TELO2, ATF4 and HIF1A (PubMed:24809345). Target proteins are preferentially recognized via a LXXLAP motif (By similarity). Cellular oxygen sensor that catalyzes, under normoxic conditions, the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins (By similarity). Hydroxylates a specific proline found in each of the oxygen-dependent degradation (ODD) domains (N-terminal, NODD, and C-terminal, CODD) of HIF1A (By similarity). Also hydroxylates HIF2A (By similarity). Has a preference for the CODD site for both HIF1A and HIF2A (By similarity). Hydroxylation on the NODD site by EGLN3 appears to require prior hydroxylation on the CODD site (By similarity). Hydroxylated HIFs are then targeted for proteasomal degradation via the von Hippel-Lindau ubiquitination complex (By similarity). Under hypoxic conditions, the hydroxylation reaction is attenuated allowing HIFs to escape degradation resulting in their translocation to the nucleus, heterodimerization with HIF1B, and increased expression of hypoxia-inducible genes (By similarity). ELGN3 is the most important isozyme in limiting physiological activation of HIFs (particularly HIF2A) in hypoxia (By

Target Details

similarity). Also hydroxylates PKM in hypoxia, limiting glycolysis (By similarity). Under normoxia, hydroxylates and regulates the stability of ADRB2. Regulator of cardiomyocyte and neuronal apoptosis (By similarity). In cardiomyocytes, inhibits the anti-apoptotic effect of BCL2 by disrupting the BAX-BCL2 complex (By similarity). In neurons, has a NGF-induced proapoptotic effect, probably through regulating CASP3 activity (By similarity). Also essential for hypoxic regulation of neutrophilic inflammation (PubMed:21317538). Plays a crucial role in DNA damage response (DDR) by hydroxylating TEL2, promoting its interaction with ATR which is required for activation of the ATR/CHK1/p53 pathway (By similarity). Also mediates hydroxylation of ATF4, leading to decreased protein stability of ATF4 (PubMed:24809345). {ECO:0000250|UniProtKB:Q9H6Z9, ECO:0000269|PubMed:21317538, ECO:0000269|PubMed:24809345}.

Molecular Weight: 27.3 kDa

UniProt: [Q91UZ4](#)

Pathways: [Positive Regulation of Endopeptidase Activity](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months