

[Go to Product page](#)

Datasheet for ABIN7565027

**RENT1/UPF1 Protein (AA 1-1124) (His tag)**

## Overview

Quantity:	1 mg
Target:	RENT1/UPF1 (UPF1)
Protein Characteristics:	AA 1-1124
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This RENT1/UPF1 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

## Product Details

Purpose:	Custom-made recombinat Upf1 Protein expressed in mammalien cells.
Sequence:	MSVEAYGPSS QLTFLDTEE AELLGADTQG SEFEFTDFTL PSQTQTTPPGG PGGAGGPGGA GAGGAAGQLD AQVGPEGILQ NGAVDDSVAK TSQLLAELNF EEDEEDTYT KDLPVHACSY CGIHDPACVV YCNTSKKWFC NGRGNTSGSH IVNHLVRAKC KEVTLHKDGP LGETVLECYN CGCRNVFLG FIPAKADSVV VLLCRQPCAS QSSLKDINWD SSQWQPLIQD RCFLSWLVKI PSEQEQLRAR QITAQQINKL EELWKENPSA TLEDLEKPGV DEEPQHVLLR YEDAYQYQNI FGPLVKLEAD YDKKLESQT QDNITVRWDL GLNKKRIAFF TLPKTDGNE DLVIIWLRDM RLMQGDEICL RYKGD LAPLW KGIGHVIKVP DNYGDEIAIE LRSSVGAPVE VTHNFQVDFV WKSTSFDRMQ SALKTFVAVDE TSVSGYIYHK LLGHEVEDVV IKCQLPKRFT AQGLPDLNHS QVYAVKTVLQ RPLSLIQGPP GTGKTVTSAT IVYHLARQGN GPVLVCAPSN IAVDQLTEKI HQTGLKVVRL CAKSREIDS PVSFLALHNQ IRNMDSMPPEL QKLQQLKDET GELSSADEKR YRALKRTER ELLMNADVIC CTCVGAGDPR LAKMQFRSIL IDESTQATEP ECMVPVWLGA

## Product Details

---

KQLILVGDHC QLGPVVMCKK AAKAGLSQSL FERLVVLGIR PIRLQVQYRM HPALSAFPSN  
IFYEGSLQNG VTAADRVKKG FDFQWPQPK PMFFYVTQGG EEIASSGTSY LNRTEANVE  
KITTKLLKAG AKPDQIGIIT PYEGQRSYLV QYMQFSGSLH TKLYQEVEIA SVDAFQGREK  
DFIILSCVRA NEHQGIGFLN DPRRLNVALT RARYGVIIVG NPKALSKQPL WNHLLSYYKE  
QKALVEGPLN NLRESLMQFS KPRKLVNTVN PGARFMTTAM YDAREAIIPG SVYDRSSQGR  
PSNMYFQTHD QISMISAGPS HVAAMNIPIP FNLVMPMPMP PGYFGQANGP AAGRGTPTKTK  
TGRGGRQKNR FGLPGSQTT LPNSQASQDV ASQPFSQ GAL TQGYVSMSQP SQMSQPGLSQ  
PELSQDSYLG DEFKSQIDVA LSQDSTYQGE RAYQHGGVTG LSQY **Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

---

### Characteristics:

#### Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

---

### Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

---

### Grade:

custom-made

---

## Target Details

---

### Target:

RENT1/UPF1 (UPF1)

---

### Alternative Name:

Upf1 ([UPF1 Products](#))

---

### Background:

Regulator of nonsense transcripts 1 (EC 3.6.4.12) (EC 3.6.4.13) (ATP-dependent helicase RENT1) (Nonsense mRNA reducing factor 1) (NORF1) (Up-frameshift suppressor 1 homolog)

## Target Details

---

(mUpf1),FUNCTION: RNA-dependent helicase required for nonsense-mediated decay (NMD) of aberrant mRNAs containing premature stop codons and modulates the expression level of normal mRNAs (By similarity). Is recruited to mRNAs upon translation termination and undergoes a cycle of phosphorylation and dephosphorylation, its phosphorylation appears to be a key step in NMD (By similarity). Recruited by release factors to stalled ribosomes together with the SMG1C protein kinase complex to form the transient SURF (SMG1-UPF1-eRF1-eRF3) complex (By similarity). In EJC-dependent NMD, the SURF complex associates with the exon junction complex (EJC) (located 50-55 or more nucleotides downstream from the termination codon) through UPF2 and allows the formation of an UPF1-UPF2-UPF3 surveillance complex which is believed to activate NMD (By similarity). Phosphorylated UPF1 is recognized by EST1B/SMG5, SMG6 and SMG7 which are thought to provide a link to the mRNA degradation machinery involving exonucleolytic and endonucleolytic pathways, and to serve as adapters to protein phosphatase 2A (PP2A), thereby triggering UPF1 dephosphorylation and allowing the recycling of NMD factors (By similarity). UPF1 can also activate NMD without UPF2 or UPF3, and in the absence of the NMD-enhancing downstream EJC indicative for alternative NMD pathways (By similarity). Plays a role in replication-dependent histone mRNA degradation at the end of phase S, the function is independent of UPF2 (By similarity). For the recognition of premature termination codons (PTC) and initiation of NMD a competitive interaction between UPF1 and PABPC1 with the ribosome-bound release factors is proposed (By similarity). The ATPase activity of UPF1 is required for disassembly of mRNPs undergoing NMD (By similarity). Together with UPF2 and dependent on TDRD6, mediates the degradation of mRNA harboring long 3'UTR by inducing the NMD machinery (PubMed:27149095). Also capable of unwinding double-stranded DNA and translocating on single-stranded DNA (By similarity). {ECO:0000250|UniProtKB:Q92900, ECO:0000269|PubMed:27149095}.

---

Molecular Weight: 124.0 kDa

---

UniProt: [Q9EPU0](#)

---

Pathways: [SARS-CoV-2 Protein Interactome](#)

## Application Details

---

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

---

Restrictions: For Research Use only

## Handling

---

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months