

Datasheet for ABIN7565183  
**CRY2 Protein (AA 1-592) (His tag)**



[Go to Product page](#)

## Overview

Quantity:	1 mg
Target:	CRY2
Protein Characteristics:	AA 1-592
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CRY2 protein is labelled with His tag.

## Product Details

Purpose:	Custom-made recombinant Cry2 Protein expressed in mammalian cells.
Sequence:	<p>MAAAAVVAAT VPAQSMGADG ASSVHWFRKG LRLHDNPALL AAVRGARCVR CVYILDPWFA          ASSSVGINRW RFLQSLLEDL DTSRLRKLNSR LFVVRGQPAD VFPRLFKEWG VTRLTFEYDS          EPFGKERDAA IMKMAKEAGV EVTENSHTL YDLDRIIELN GQKPPLTYKR FQALISRMEL          PKKPAVAVSS QQMESCR AEI QENHDDTYGV PSLEELGFPT EGLGPAVWQG GETEALARLD          KHLERKAWVA NYERPRMNAN SLLASPTGLS PYLRFGLCLSC RLFYYRLWDL YKKVKRNSTP          PLSLFGQLLW REFFYTAATN NPRFDRMEGN PICIQIPWDR NPEALAKWAE GKTGFPWIDA          IMTQLRQEGW IHHLARHAVA CFLTRGDLWV SWESGVRVFD ELLLDADFSV NAGSWMWLSC          SAFFQQFFHC YCPVGFGRRT DPSGDYIRRY LPKLGFPSPR YIYEPWNAPE SVQKAAKCII          GVDYPRPIVN HAETSRLNIE RMKQIQQLS RYRGLCLLAS VPSCVEDLSH PVAEPGSSQA          GSI SNTGPRA LSSGPASPKR KLEAAEPPG EELTKRARVT EMPTQEPASK DS <b>Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a</b></p>

## Product Details

---

### **special request, please contact us.**

**Specificity:** If you are looking for a specific domain and are interested in a partial protein or a different isoform, please contact us regarding an individual offer.

**Characteristics:**

**Key Benefits:**

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

**Purity:** > 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)

**Grade:** custom-made

## Target Details

---

**Target:** CRY2

**Alternative Name:** Cry2 ([CRY2 Products](#))

**Background:** Cryptochrome-2,FUNCTION: Transcriptional repressor which forms a core component of the circadian clock. The circadian clock, an internal time-keeping system, regulates various physiological processes through the generation of approximately 24 hour circadian rhythms in gene expression, which are translated into rhythms in metabolism and behavior. It is derived from the Latin roots 'circa' (about) and 'diem' (day) and acts as an important regulator of a wide array of physiological functions including metabolism, sleep, body temperature, blood pressure, endocrine, immune, cardiovascular, and renal function. Consists of two major components: the central clock, residing in the suprachiasmatic nucleus (SCN) of the brain, and the peripheral clocks that are present in nearly every tissue and organ system. Both the central and peripheral clocks can be reset by environmental cues, also known as Zeitgebers (German for 'timegivers').

The predominant Zeitgeber for the central clock is light, which is sensed by retina and signals directly to the SCN. The central clock entrains the peripheral clocks through neuronal and hormonal signals, body temperature and feeding-related cues, aligning all clocks with the external light/dark cycle. Circadian rhythms allow an organism to achieve temporal homeostasis with its environment at the molecular level by regulating gene expression to create a peak of protein expression once every 24 hours to control when a particular physiological process is most active with respect to the solar day. Transcription and translation of core clock components (CLOCK, NPAS2, BMAL1, BMAL2, PER1, PER2, PER3, CRY1 and CRY2) plays a critical role in rhythm generation, whereas delays imposed by post-translational modifications (PTMs) are important for determining the period ( $\tau$ ) of the rhythms ( $\tau$  refers to the period of a rhythm and is the length, in time, of one complete cycle). A diurnal rhythm is synchronized with the day/night cycle, while the ultradian and infradian rhythms have a period shorter and longer than 24 hours, respectively. Disruptions in the circadian rhythms contribute to the pathology of cardiovascular diseases, cancer, metabolic syndromes and aging. A transcription/translation feedback loop (TTFL) forms the core of the molecular circadian clock mechanism. Transcription factors, CLOCK or NPAS2 and BMAL1 or BMAL2, form the positive limb of the feedback loop, act in the form of a heterodimer and activate the transcription of core clock genes and clock-controlled genes (involved in key metabolic processes), harboring E-box elements (5'-CACGTG-3') within their promoters. The core clock genes: PER1/2/3 and CRY1/2 which are transcriptional repressors form the negative limb of the feedback loop and interact with the CLOCK|NPAS2-BMAL1|BMAL2 heterodimer inhibiting its activity and thereby negatively regulating their own expression. This heterodimer also activates nuclear receptors NR1D1/2 and RORA/B/G, which form a second feedback loop and which activate and repress BMAL1 transcription, respectively. CRY1 and CRY2 have redundant functions but also differential and selective contributions at least in defining the pace of the SCN circadian clock and its circadian transcriptional outputs. Less potent transcriptional repressor in cerebellum and liver than CRY1, though less effective in lengthening the period of the SCN oscillator. Seems to play a critical role in tuning SCN circadian period by opposing the action of CRY1. With CRY1, dispensable for circadian rhythm generation but necessary for the development of intercellular networks for rhythm synchrony. May mediate circadian regulation of cAMP signaling and gluconeogenesis by blocking glucagon-mediated increases in intracellular cAMP concentrations and in CREB1 phosphorylation. Besides its role in the maintenance of the circadian clock, is also involved in the regulation of other processes. Plays a key role in glucose and lipid metabolism modulation, in part, through the transcriptional regulation of genes involved in these pathways, such as LEP or ACSL4. Represses glucocorticoid receptor NR3C1/GR-induced transcriptional activity by binding to glucocorticoid response elements (GREs). Represses the CLOCK-BMAL1 induced

## Target Details

---

transcription of BHLHE40/DEC1 and NAMPT. Represses PPARD and its target genes in the skeletal muscle and limits exercise capacity (PubMed:28683290). Represses the transcriptional activity of NR1I2 (PubMed:28751364). {ECO:0000269|PubMed:10428031, ECO:0000269|PubMed:16628007, ECO:0000269|PubMed:17310242, ECO:0000269|PubMed:19299583, ECO:0000269|PubMed:20852621, ECO:0000269|PubMed:22170608, ECO:0000269|PubMed:23531614, ECO:0000269|PubMed:23575670, ECO:0000269|PubMed:23616524, ECO:0000269|PubMed:24154698, ECO:0000269|PubMed:28683290, ECO:0000269|PubMed:28751364}.

---

Molecular Weight: 66.9 kDa

---

UniProt: [Q9R194](#)

---

Pathways: [Response to Water Deprivation, Protein targeting to Nucleus](#)

## Application Details

---

Application Notes: We expect the protein to work for functional studies. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Buffer: The buffer composition is at the discretion of the manufacturer.

---

Handling Advice: Avoid repeated freeze-thaw cycles.

---

Storage: -80 °C

---

Storage Comment: Store at -80°C.

---

Expiry Date: 12 months