

Datasheet for ABIN7565743 anti-Avidin antibody (FITC)



Overview

Quantity:	25 μL
Target:	Avidin (AVD)
Reactivity:	Chicken
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Avidin antibody is conjugated to FITC
Application:	Western Blotting (WB), Flow Cytometry (FACS), Dot Blot (DB), Fluorescence Microscopy (FM), FLISA

Product Details

Purpose:	Avidin Antibody Fluorescein Conjugated
Immunogen:	Immunogen: Avidin (Hen Egg White) Immunogen Type: Native Protein
Cross-Reactivity (Details):	Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-fluorescein, anti-Rabbit Serum and Avidin.
Characteristics:	Synonyms: rabbit anti-Avidin Antibody fluorescein Conjugation, FITC conjugated rabbit anti-Avidin antibody, Avidin fluorescein, Anti-Avidin FITC Antibody, egg ehite
Purification:	Anti-Avidin Antibody Fluorescein Conjugated is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above.

Product Details Sterility: Sterile filtered Labeling Ratio: 3.0 Target Details Target: Avidin (AVD) Alternative Name: **AVIDIN (AVD Products)** Background: Background: Avidin is a glycoprotein with a molecular weight of approximately 62.4 kDa. Avidin is a biotin binding protein that shows high sequence homology in birds, reptiles and amphibians. Hen egg white avidin is a tetrameric protein composed of four identical subunits, each with the ability to bind biotin with high affinity and specificity (Kd \sim 1015 M). In biotechnology, the functional consequence of tetrameric biotin binding is signal amplification. Biotin-avidin bridging is a great way to increase signal strength while maintaining specificity. The sequence of avidin only shows 30 % homology with streptavidin, and anti-avidin and antistreptavidin antibodies are not immunologically cross reactive. UniProt: P02701 **Application Details Application Notes:** Application Note: Rabbit Anti-Avidin Fluorescein conjugate has been tested by dot blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. Flow Cytometry Dilution: User Optimized Western Blot Dilution: >1:1,000 FLISA Dilution: >1:1,000 IF Microscopy Dilution: User Optimized Other: User Optimized Restrictions: For Research Use only Handling

Format: Concentration: 1.0 mg/mL Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Preservative: 0.01 % (w/v) Sodium Azide Preservative: Sodium azide

Handling

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 μ L). To minimize loss of volume dilute 1:10 by adding 225 μ L of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Expiry Date:	12 months