

Datasheet for ABIN7565836

anti-RFP antibody



Overview

| Quantity: | 25 μL |
|--------------|---|
| Target: | RFP |
| Reactivity: | Discosoma |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This RFP antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunoprecipitation (IP), Flow Cytometry (FACS), Fluorescence Microscopy (FM) |

Product Details

| Purpose: | RFP Antibody |
|-----------------------------|---|
| Immunogen: | Immunogen: The immunogen is a Red Fluorescent Protein (RFP) fusion protein corresponding to the full-length amino acid sequence (234aa) derived from the mushroom anemone Discosoma. Immunogen Type: Recombinant Protein |
| Clone: | 8E5-G7 |
| Isotype: | IgG2a kappa |
| Cross-Reactivity (Details): | Expect reactivity against RFP and its variants: mCherry, tdTomato, mBanana, mOrange, mPlum, mOrange and mStrawberry. |
| Characteristics: | Synonyms: mouse anti-RFP Antibody, DsRed, rDsRed, Discosoma sp. Red Fluorescent Protein, Red fluorescent protein drFP583 |

| Anti-RFP Monoclonal Antibody was purified from concentrated tissue culture supernate by |
|---|
| Protein A chromatography. |
| Sterile filtered |
| |
| RFP |
| DsRed (RFP Products) |
| Background: Antibodies to RFP (Discosoma spp.) are intended for use in immunological assays |
| including ELISA, Western blotting, immunohistochemistry, and flow cytometry. RFP Proteins are |
| useful markers for imagining protein localization, monitoring physiological processes, and |
| detecting transgenic expression. Rockland's anti-RFP antibody can be used to detect native |
| RFP and RFP variants. |
| Q9U6Y8 |
| |
| Application Note: Monoclonal anti-RFP is designed to detect RFP and its variants. This antibody |
| has been tested by ELISA and Western blot, and is suitable for use in FISH, IF, IHC, and |
| multiplex assays based on published references. This antibody can be used to detect RFP by |
| ELISA (sandwich or capture) for the direct binding of antigen. Optimal titers for applications |
| should be determined by the researcher. Flow Cytometry Dilution: User Optimized Western Blot |
| Dilution: 1:1,000 - 1:10,000 Immunoprecipitation Dilution: User Optimized ELISA Dilution: |
| 1:75,000 - 1:150,000 IF Microscopy Dilution: User Optimized |
| For Research Use only |
| |
| Liquid |
| Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Stabilizer: None |
| Preservative: 0.01 % (w/v) Sodium Azide |
| Sodium azide |
| This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which |
| |

should be handled by trained staff only.

Handling

| Storage: | -20 °C |
|------------------|--|
| Storage Comment: | Store Anti-RFP at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 μ L). To minimize loss of volume dilute 1:10 by adding 225 μ L of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing. |
| Expiry Date: | 12 months |