

Datasheet for ABIN7566036 **BMP2 Protein (AA 289-396) (His tag)**



Go to Product page

	er		

Quantity:	10 μg
Target:	BMP2
Protein Characteristics:	AA 289-396
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This BMP2 protein is labelled with His tag.

Product Details

Purpose:	BMP-2 (human) (rec.) (His)
Cross-Reactivity:	Human
Characteristics:	Human BMP-2 (aa 289-396) is fused at the C-terminus to a His-tag.
Purity:	>95 % (SDS-PAGE)
Sterility:	Sterile filtered
Endotoxin Level:	<0.1EU/µg protein (LAL test, Lonza).

Target Details

Target:	BMP2
Alternative Name:	BMP-2 (BMP2 Products)
Background:	BMP2, Bone Morphogenetic Protein 2

Bone morphogenetic protein 2 (BMP-2) is a member of the BMP subgroup of the TGF-beta superfamily. It plays a dominant role in embryonic dorsalventral patterning, organogenesis, limb bud formation and bone formation and regeneration. BMP-2 signals through heterodimeric complexes composed of a type I receptor (Activin RI, BMP-RIA or BMP-RIB) and a type II receptor (BMP-RII or Activin RIIB). BMP-2 induces chondrocyte proliferation, endochondral bone formation, longitudinal bone growth and bone and cartilage repair. It induces ectopic bone formation or calcification by promoting osteogenic and chondrogenic differentiation in mesenchymal cells, stem cells and vascular smooth muscle cells. It also promotes the maintenance and repair of colonic epithelium, suppresses neuronal dopamine synthesis and release, induces apoptosis in medulloblastoma cells and is required for cardiac contractility.

NCBI Accession:

NP_001191

Pathways:

Regulation of Hormone Metabolic Process, Regulation of Hormone Biosynthetic Process,
Regulation of Muscle Cell Differentiation, Growth Factor Binding, Positive Regulation of fat Cell
Differentiation

Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized		
Buffer:	Lyophilized from 0.2µm-filtered solution in PBS.		
Handling Advice:	Avoid freeze/thaw cycles.Centrifuge lyophilized vial before opening and reconstitution.		
Storage:	4 °C,-20 °C		
Storage Comment:	Short Term Storage: +4°C		
	Long Term Storage: -20°C		
	Use & Stability: Stable for at least 1 year after receipt when stored at -20°C. Working aliquots		
	are stable for up to 3 months when stored at -20°C.		