

Datasheet for ABIN7566068

TNFSF4 Protein (AA 51-183) (His tag)



| () | ve | r\/i | Δ | ۱۸/ |
|---------|-----|------|---|-----|
| \circ | V C | 1 V | | v v |

| Quantity: | 50 µg |
|-------------------------------|---|
| Target: | TNFSF4 |
| Protein Characteristics: | AA 51-183 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Biological Activity: | Active |
| Purification tag / Conjugate: | This TNFSF4 protein is labelled with His tag. |
| | |

Product Details

| Purpose: | CD252 [OX40L] (human) (rec.) (His) |
|------------------------------|--|
| Cross-Reactivity: | Human, Mouse |
| Characteristics: | The extracellular domain of human CD252 (aa 51-183) is fused at the C-terminus to a His-tag. |
| Purity: | >95 % (SDS-PAGE) |
| Sterility: | Sterile filtered |
| Endotoxin Level: | <0.01EU/µg purified protein (LAL test, Lonza). |
| Biological Activity Comment: | Measured by its binding ability in a functional ELISA. |

Target Details

Target: TNFSF4

Target Details

| Alternative Name: | CD252 [OX40L] (TNFSF4 Products) | |
|-------------------|---|--|
| Background: | TNFSF4 OX40L [CD252, TNFSF4] is a membrane-expressed cytokine that belongs to the TNF ligand family. It acts as costimulator through interaction with its ligand OX40 on T cells, stimulating T cell activation, proliferation and cytokine production. OX40L is expressed on antigen presenting cells including B cells, dendritic cells, mast cells and endothelium. | |
| Molecular Weight: | ~27kDa (SDS-PAGE) | |
| NCBI Accession: | NP_003317 | |
| Pathways: | Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, Activated T Cell Proliferation, Cancer Immune Checkpoints | |

Application Details

|--|--|

Handling

| Format: | Lyophilized | |
|------------------|--|--|
| Buffer: | Lyophilized from 0.2µm-filtered solution in PBS. | |
| Handling Advice: | Avoid freeze/thaw cycles. Centrifuge lyophilized vial before opening and reconstitution. PBS containing at least 0.1 % BSA should be used for further dilutions. | |
| Storage: | 4 °C,-20 °C | |
| Storage Comment: | Short Term Storage: +4°C Long Term Storage: -20°C Use & Stability: Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C. | |