

Datasheet for ABIN7566186

IL12 Protein (AA 1-335, AA 23-215) (Fc Tag)[Go to Product page](#)

Overview

Quantity:	25 µg
Target:	IL12
Protein Characteristics:	AA 1-335, AA 23-215
Origin:	Mouse
Source:	CHO Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL12 protein is labelled with Fc Tag.

Product Details

Purpose:	IL-12 (mouse):Fc (human) (rec.)
Cross-Reactivity:	Mouse
Characteristics:	The mouse IL-12 complex composed of the p40 subunit (aa 1-335) and the p35 subunit (aa 23-215) is fused through a polypeptide linker to the Fc region of human IgG1.
Purity:	>98 % (SDS-PAGE)
Sterility:	Sterile filtered
Endotoxin Level:	<0.06EU/µg protein (LAL test, Lonza).

Target Details

Target:	IL12
Alternative Name:	IL-12 (IL12 Products)

Target Details

Background:	IL12-p70, CLMF, Cytotoxic Lymphocyte Maturation Factor, NKSF, Natural Killer Cell Stimulatory Factor, TSF, T cell Stimulating Factor Interleukin-12 (IL-12), also known as Natural Killer Cell Stimulatory Factor (NKSF) or Cytotoxic Lymphocyte Maturation Factor (CLMF), is a heterodimeric pleiotropic cytokine made up of a 40 kDa (p40) subunit and a 35 kDa (p35) subunit. IL-12 is produced by macrophages and B lymphocytes and has been shown to have multiple effects on T cells and Natural Killer (NK) cells. Some of these IL-12 activities include the induction of IFN-gamma and TNF in resting and activated T and NK cells, the enhancement of cytotoxic activity of resting NK and T cells, the stimulation of resting T cell proliferation in the presence of a comitogen, and the enhancement of NK cell proliferation. Current evidence indicates that IL-12 is a key mediator of cellular immunity and induces the differentiation of Th1 cells from precursor T helper cells. Based on its activities, it has been suggested that IL-12 may have therapeutic potential as a vaccine adjuvant that promotes cellular immunity and as an antitumor and anti viral agent.
Molecular Weight:	~100-110kDa (SDS-PAGE, reducing conditions)
NCBI Accession:	NP_032377
Pathways:	JAK-STAT Signaling , TLR Signaling , Cellular Response to Molecule of Bacterial Origin , Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process , Activated T Cell Proliferation , Cancer Immune Checkpoints , Inflammasome

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Buffer:	Lyophilized from 0.2µm-filtered solution in PBS.
Handling Advice:	Avoid freeze/thaw cycles. Centrifuge lyophilized vial before opening and reconstitution.
Storage:	4 °C, -20 °C
Storage Comment:	Short Term Storage: +4°C Long Term Storage: -20°C Use & Stability: Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.