

Datasheet for ABIN7566268

Interleukin 35 Protein (IL35) (AA 23-215, AA 23-228) (Fc Tag)[Go to Product page](#)

Overview

Quantity:	5 µg
Target:	Interleukin 35 (IL35)
Protein Characteristics:	AA 23-215, AA 23-228
Origin:	Mouse
Source:	CHO Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This Interleukin 35 protein is labelled with Fc Tag.

Product Details

Purpose:	IL-35 (mouse):Fc (human) (rec.)
Cross-Reactivity:	Mouse
Characteristics:	A soluble dimeric fusion protein consisting of the extracellular domain of mouse IL12a subunit (aa 23-215) is fused to the Fc region of human IgG1, and the mouse Ebi3 subunit (aa 23-228) linked to IL12a by disulfide bonds.
Purity:	>95 % (SDS-PAGE)
Sterility:	Sterile filtered
Endotoxin Level:	<1EU/mg protein (LAL test, Lonza).
Biological Activity Comment:	Bioactivity was measured in a cell proliferation assay of Con A activated mouse splenocytes.

Target Details

Target:	Interleukin 35 (IL35)
Alternative Name:	IL-35 (IL35 Products)
Background:	<p>Interleukin-35</p> <p>Interleukin-35 (IL-35) is a novel IL-12 family cytokine produced by regulatory T cells (Treg) but not by resting or activated effector T cells (Teff). IL-35 is a heterodimeric protein composed of EBI3 (Epstein-Barr-Virus-induced gene 3) and IL-12a (p35). EBI3 is a downstream target of Foxp3, a transcription factor required for Treg-cell development and function, and thus Treg-cell restriction of IL35 occurs. Regulatory T cells are essential for maintaining self tolerance and preventing autoimmunity, and IL-35 is identified as a molecule that mediates the immune suppression function of Treg-cell. As an inhibitory cytokine, IL-35 induces proliferation of Treg-cell populations but suppresses Th17 cell development. Studies in mice show the absence of either IL-35 chain from Treg-cell reduces the cells' ability to suppress inflammation using an experimental model for inflammatory bowel disease. IL-35 is suggested as a potential target of immunotherapy. Recently, insufficient IL-35 levels were shown to play a pivotal role in the development of type 1 diabetes (T1D) and autoimmune diseases.</p>
Molecular Weight:	~30-60kDa
NCBI Accession:	NP_032377

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Buffer:	Lyophilized from 0.2µm-filtered solution in PBS.
Handling Advice:	Avoid freeze/thaw cycles. Centrifuge lyophilized vial before opening and reconstitution.
Storage:	4 °C, -20 °C
Storage Comment:	<p>Short Term Storage: +4°C</p> <p>Long Term Storage: -20°C</p> <p>Use & Stability: Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.</p>