

Datasheet for ABIN7566281 **IL1F10 Protein (AA 3-152) (DYKDDDDK Tag)**



Go to Product page

Overviev	

Quantity:	3 x 10 μg
Target:	IL1F10
Protein Characteristics:	AA 3-152
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL1F10 protein is labelled with DYKDDDDK Tag.

Product Details

Purpose:	IL-38 (aa 3-152) (mouse) (rec.) (FLAG)
Cross-Reactivity:	Mouse
Characteristics:	Mouse IL-38 [IL-1F10] (aa 3-152) is fused at the C-terminus to a DYKDDDDK Tag.
Purity:	>95 % (SDS-PAGE)
Endotoxin Level:	<0.01EU/µg purified protein (LAL test, Lonza).

Target Details

Target:	IL1F10
Alternative Name:	IL-38 (IL1F10 Products)
Background:	Interleukin-38, Interleukin-1 Family Member 10, IL-1F10
	IL-38 (IL-1F10) mRNA is expressed in heart, placenta, fetal liver, spleen, thymus and tonsil. The

Target Details

expression in a variety of immune tissues and similarity to IL-1Ra suggest a role of IL-1F10 in the inflammatory response. It has been reported that removal of the N-terminus domain of the interleukins of the IL-1F family such as IL-1F5 / 6 / 8 or 9 (also called IL-36Ra, IL-36alpha, IL-36beta or IL-36gamma) is important to increase their biological activity. Recently it has been shown that IL-38 is N-terminally processed and secreted by apoptotic cells. Released processed IL-38 (20-152) binds to the receptor Interleukin-1 receptor accessory protein-like 1 (IL1RAPL1, TIGIRR-2) at the surface of macrophages. Processed IL-38-activated IL1RAPL1 reduces the production of IL-6 leading to inflammation attenuation.

Molecular Weight:

~20kDa

Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	0.1 mg/mL after reconstitution.
Concentration:	0.1 mg/mL
Buffer:	Contains PBS + 0.5 % Trehalose + 0.02 % CHAPS.
Handling Advice:	After reconstitution, prepare aliquots and store at -20 °C. Avoid freeze/thaw cycles. Centrifuge lyophilized vial before opening and reconstitution. PBS containing at least 0.1 % BSA should be used for further dilutions.
Storage:	4 °C,-20 °C
Storage Comment:	Short Term Storage: +4°C Long Term Storage: -20°C Use & Stability: Stable for at least 6 months after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.