

Datasheet for ABIN7566282 IL1F10 Protein (AA 1-152) (His tag)



Overview	
Quantity:	10 µg
Target:	IL1F10
Protein Characteristics:	AA 1-152
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL1F10 protein is labelled with His tag.
Product Details	
Purpose:	IL-38 (human) (rec.) (His)
Cross-Reactivity:	Human
Characteristics:	Human IL-38 (aa 1-152) is fused at the C-terminus to a His-tag.
Purity:	>95 % (SDS-PAGE)
Endotoxin Level:	<1EU/µg purified protein (LAL test, Lonza).

Target Details

Target:	IL1F10
Alternative Name:	IL-38 (IL1F10 Products)
Background:	Interleukin-38, Interleukin-1 Family Member 10 theta, IL-1F10, FIL1 theta, FKSG75, IL-1HY2, IL1- theta, Interleukin-1 HY2, Interleukin-1 Receptor Antagonist FKSG75

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7566282 | 04/22/2025 | Copyright antibodies-online. All rights reserved.

	IL-38 (IL-1F10) mRNA is expressed in heart, placenta, fetal liver, spleen, thymus and tonsil. The expression in a variety of immune tissues and similarity to IL-1Ra suggest a role of IL-1F10 in the inflammatory response. It has been reported that removal of the N-terminus domain of the interleukins of the IL-1F family such as IL-1F5 / 6 / 8 or 9 (also called IL-36Ra, IL-36alpha, IL-36beta or IL-36gamma) is important to increase their biological activity. Recently it has been shown that IL-38 is N-terminally processed and secreted by apoptotic cells. Released processed IL-38 (20-152) binds to the receptor Interleukin-1 receptor accessory protein-like 1 (IL1RAPL1, TIGIRR-2) at the surface of macrophages. Processed IL-38-activated IL1RAPL1 reduces the production of IL-6 leading to inflammation attenuation.
Molecular Weight:	~18kDa (SDS-PAGE)
UniProt:	Q8WWZ1
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	After reconstitution:for 10 µg size: 0.1 mg/mL
Reconstitution: Concentration:	After reconstitution:for 10 µg size: 0.1 mg/mL Lot specific
Concentration:	Lot specific
Concentration: Buffer:	Lot specific In 50 mM TRIS-Cl, pH 8.0, containing 150 mM NaCl.