

Datasheet for ABIN7566307 **IL-7 Protein (AA 26-177) (Fc Tag)**



Overview

| Quantity: | 50 μg |
|-------------------------------|--------------------------------------------|
| Target: | IL-7 (IL7) |
| Protein Characteristics: | AA 26-177 |
| Origin: | Human |
| Source: | CHO Cells |
| Protein Type: | Recombinant |
| Biological Activity: | Active |
| Purification tag / Conjugate: | This IL-7 protein is labelled with Fc Tag. |

Product Details

| Purpose: | IL-7 (human):Fc (human) (rec.) (non-lytic) |
|------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Cross-Reactivity: | Human |
| Characteristics: | The extracellular domain of human IL-7 (aa 26-177) is fused to the N-terminus of the Fc portion of a mutant human IgG1. |
| Purity: | >98 % (SDS-PAGE) |
| Sterility: | Sterile filtered |
| Endotoxin Level: | <1EU/mg protein (LAL test, Lonza). |
| Biological Activity Comment: | Measured in a cell proliferation assay using antibody against CD3-activated human peripheral blood lymphocytes (PBL). The ED50 for this effect is typically 10-100ng/ml. |

Target Details

| Target: | IL-7 (IL7) |
|---------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Alternative Name: | IL-7 (IL7 Products) |
| Background: | Interleukin-7 Interleukin-7 (IL-7), is a hematopoietic growth factor and a member of the IL-7/IL-9 family. IL-7 stimulates the proliferation of lymphoid progenitors and is secreted by stromal cells in the bone marrow and thymus. IL-7 is also important for proliferation during certain stages of B cell maturation. IL-7 and the hepatocyte growth factor (HGF) form a heterodimer that functions as a pre-pro-B cell growth-stimulating factor. It is found to be a cofactor for V(D)J rearrangement of the T cell receptor beta (TCRbeta) during early T cell development. IL-7 can be produced locally by intestinal epithelial and epithelial goblet cells and may serve as a regulatory factor for intestinal mucosal lymphocytes. |
| Molecular Weight: | ~50-60kDa |
| Pathways: | JAK-STAT Signaling |
| Application Details | |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Buffer: | Lyophilized from 0.2µm-filtered solution in PBS. |
| Handling Advice: | Avoid freeze/thaw cycles.Centrifuge lyophilized vial before opening and reconstitution. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Short Term Storage: +4°C Long Term Storage: -20°C Use & Stability: Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C. |