

## Datasheet for ABIN7566336

## SARS-CoV-2 Spike S1 Protein (AA 319-541) (His tag, Biotin)



Go to Product page

_						
	V	$\triangle$	r۱	/1	$\triangle$	Λ/
	' V '		ΙV			v v

Quantity:	50 µg	
Target:	SARS-CoV-2 Spike S1	
Protein Characteristics:	AA 319-541	
Origin:	SARS Coronavirus-2 (SARS-CoV-2)	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Biological Activity:	Active	
Purification tag / Conjugate:	This SARS-CoV-2 Spike S1 protein is labelled with His tag, Biotin.	
Product Details		
_		
Purpose:	SARS-CoV-2 Spike Protein S1 (RBD) (rec.) (His) (Biotin)	
Purpose:  Cross-Reactivity:	SARS-CoV-2 Spike Protein S1 (RBD) (rec.) (His) (Biotin)  Human	
Cross-Reactivity:	Human	
Cross-Reactivity:	Human  Receptor-binding domain (RBD) of SARS-CoV-2 Spike protein S1 (aa 319-541) is fused at the C-	
Cross-Reactivity: Characteristics:	Human  Receptor-binding domain (RBD) of SARS-CoV-2 Spike protein S1 (aa 319-541) is fused at the C-terminus to a His-tag.	
Cross-Reactivity: Characteristics: Purity:	Human  Receptor-binding domain (RBD) of SARS-CoV-2 Spike protein S1 (aa 319-541) is fused at the C-terminus to a His-tag.  >95 % (SDS-PAGE)	
Cross-Reactivity:  Characteristics:  Purity:  Endotoxin Level:	Human  Receptor-binding domain (RBD) of SARS-CoV-2 Spike protein S1 (aa 319-541) is fused at the C-terminus to a His-tag.  >95 % (SDS-PAGE)  <0.01EU/µg purified protein (LAL test).	
Cross-Reactivity: Characteristics:  Purity: Endotoxin Level: Biological Activity Comment:	Human  Receptor-binding domain (RBD) of SARS-CoV-2 Spike protein S1 (aa 319-541) is fused at the C-terminus to a His-tag.  >95 % (SDS-PAGE)  <0.01EU/µg purified protein (LAL test).	

Handling Advice:

## Abstract: SARS-CoV-2 Spike S1 Products Background: 2019-nCoV Spike Protein S1 (RBD), Spike Receptor Binding Domain SARS-CoV-2 shares 79.5 % sequence identity with SARS-CoV and is 96.2 % identical at the genome level to the bat coronavirus BatCoV RaTG133, suggesting it had originated in bats. The coronaviral genome encodes four major structural proteins: the Spike (S) protein, Nucleocapsid (N) protein, Membrane/Matrix (M) protein and the Envelope (E) protein. The SARS Envelope (E) protein contains a short palindromic transmembrane helical hairpin that seems to deform lipid bilayers, which may explain its role in viral budding and virion envelope morphogenesis. The SARS Membrane/Matrix (M) protein is one of the major structural viral proteins. It is an integral membrane protein involved in the budding of the viral particles and interacts with SARS Spike (S) protein and the Nucleocapsid (N) protein. The N protein contains two domains, both of them bind the virus RNA genome via different mechanisms. The CoV Spike (S) protein assembles as trimer and plays the most important role in viral attachment, fusion and entry. It is composed of a short intracellular tail, a transmembrane anchor and a large ectodomain that consists of a receptor binding S1 subunit (RBD domain) and a membrane-fusing S2 subunit. The S1 subunit contains a receptor binding domain (RBD), which binds to the cell surface receptor angiotensinconverting enzyme 2 (ACE2) present at the surface of epithelial cells. The SARS-CoV-2 Spike Protein S1 (RBD) (rec.) (His) is used as antigen in the Serological ELISA Kit to detect anti-SARS-CoV-2 Spike (RBD) antibodies in serum or plasma (see SARS-CoV-2 (Spike RBD) IgG Serological ELISA Kit, AG-45B-0020). This biotinylated version of SARS-CoV-2 Spike Protein S1 (RBD) (rec.) (His) forms a tetramer in the presence of streptavidin and this tetramer can be used to activate B cell memory to SARS-CoV-2 Spike protein. Molecular Weight: ~35kDa (SDS-PAGE) **Application Details** Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: 1 mg/mL after reconstitution Concentration: 1 mg/mL Buffer: Contains PBS.

After opening, prepare aliquots and store at -20 °C. Avoid freeze/thaw cycles. For maximum

## Handling

	product recovery after thawing, centrifuge the vial before opening the cap.
Storage:	4 °C,-20 °C
Storage Comment:	Short Term Storage: +4°C
	Long Term Storage: -20°C
	Use & Stability: Stable for at least 6 months after receipt when stored at -20°C. Working aliquots
	are stable for up to 3 months when stored at -20°C.