

Datasheet for ABIN7566373

HAVCR1 Protein (AA 22-212) (Fc Tag)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	HAVCR1
Protein Characteristics:	AA 22-212
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This HAVCR1 protein is labelled with Fc Tag.

Product Details

Purpose:	Tim-1 (mouse):Fc (mouse) (rec.)
Cross-Reactivity:	Mouse
Characteristics:	The extracellular domain of mouse Tim-1 (aa 22-212) is fused to the N-terminus of the Fc region of mouse IgG2a.
Purity:	>98 % (SDS-PAGE)
Sterility:	Sterile filtered
Endotoxin Level:	<5EU/mg protein (LAL test, Lonza).

Target Details

Target:	HAVCR1
Alternative Name:	Tim-1 (HAVCR1 Products)

Target Details

Background: TIM1, KIM1, TIMD1, Hepatitis A Virus Cellular Receptor 1, HAVcr-1, T Cell Immunoglobulin and Mucin Domain-containing Protein 1, T Cell Membrane Protein 1

The TIM (T cell/transmembrane, immunoglobulin and mucin) family plays a critical role in regulating immune responses, including allergy, asthma, transplant tolerance, autoimmunity and the response to viral infections. The unique structure of TIM immunoglobulin variable region domains allows highly specific recognition of phosphatidylserine (PtdSer), exposed on the surface of apoptotic cells. Tim-1 (KIM-1, HAVcr-1) is a 100 kDa, type I transmembrane glycoprotein member of the TIM family of immunoglobulin superfamily molecules. There are two cytoplasmic alternate splice forms of Tim-1. One is a long (359 aa) kidney form termed Tim-1b, and one is a short (334 aa) liver form termed Tim-1a. Tim-1, important for asthma and allergy, is preferentially expressed on T-helper 2 (Th2) cells and functions as a potent costimulatory molecule for T cell activation.

Molecular Weight: ~70kDa (SDS-PAGE)

NCBI Accession: [NP_001160104](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Buffer: Lyophilized from 0.2µm-filtered solution in PBS.

Handling Advice: Avoid freeze/thaw cycles. Centrifuge lyophilized vial before opening and reconstitution.

Storage: 4 °C, -20 °C

Storage Comment: Short Term Storage: +4°C
Long Term Storage: -20°C
Use & Stability: Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.