

Datasheet for ABIN756682 anti-SNRPE antibody (pThr401)



()	V		rV	ĺ	9	V	V
'	\mathcal{I}	٧V	<u> </u>	v	1	$\overline{}$	٧	٧

Overview		
Quantity:	100 μL	
Target:	SNRPE	
Binding Specificity:	pThr401	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SNRPE antibody is un-conjugated	
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	
Product Details		
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human B-Raf around the phosphorylation site of Thr401	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	SNRPE	

Target Details

Target Details	
Alternative Name:	B-Raf (SNRPE Products)
Background:	Synonyms: B Raf phospho T401, p-B Raf phospho T401, 94 kDa B raf protein, B raf 1, B Raf
	proto oncogene serine threonine protein kinase, BRAF 1, Braf, BRAF1, cRmil, MGC126806,
	MGC138284, Murine sarcoma viral v-raf oncogene homolog B1, Murine sarcoma viral v raf
	oncogene homolog B1, p94, RAFB 1, RAFB1, v raf murine sarcoma viral oncogene homolog B1,
	FLJ95109, BRAF_HUMAN.
	Background: The Raf kinases are important intermediates in signal transduction. Raf protein
	family members, including A Raf and B Raf, have intrinsic serine/threonine kinase activity.
	Interaction between Ras proteins and Raf proteins results in Raf-mediated phosphorylation and
	activation of MEK (also known as MAP kinase kinase). Defects in BRAF are involved in a wide
	range of cancers. B-Raf is a serine/threonine protein kinase that acts as a signal transducer
	from membrane-associated receptors to nuclear transcription factors. 1 BRAF is important for
	the regulation of cell proliferation and determination of cell fate during embryogenesis. BRAF
	acts downstream of Ras and upstream of MEK in the Ras-Raf-MEK-ERK signal transduction
	pathway, which is a conserved RAS-activated protein kinase cascade that regulates cell growth
	proliferation, and differentiation in response to growth factors, cytokines, and hormones.
Gene ID:	673
Pathways:	Ribonucleoprotein Complex Subunit Organization, Hepatitis C
Application Details	
Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months