antibodies .- online.com







anti-CDC27 antibody (pThr244)



\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	CDC27
Binding Specificity:	pThr244
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDC27 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human CDC27 around the phosphorylation site of Thr244
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Dog,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	CDC27		
---------	-------	--	--

Target Details

Alternative Name:	CDC27 (CDC27 Products)	
Background:	Synonyms: ANAPC3, Anaphase Promoting Complex 3, Anaphase promoting complex protein 3,	
	Anaphase Promoting Complex Subunit 3, Anaphase-promoting complex subunit 3, APC 3,	
	APC3, APC-3, Cdc 27, Cdc27, Cdc-27, CDC27 homolog, CDC27_HUMAN, CDC27Hs, Cell division	
	cycle 27, Cell division cycle protein 27 homolog, H NUC, H-NUC, HNUC, NUC2 homolog.	
	Background: The protein encoded by this gene shares strong similarity with Saccharomyces	
	cerevisiae protein Cdc27, and the gene product of Schizosaccharomyces pombe nuc 2. This	
	protein is a component of anaphase-promoting complex (APC), which is composed of eight	
	protein subunits and highly conserved in eucaryotic cells. APC catalyzes the formation of cyclin	
	B-ubiquitin conjugate that is responsible for the ubiquitin-mediated proteolysis of B-type cyclins	
	This protein and 3 other members of the APC complex contain the TPR (tetratricopeptide	
	repeat), a protein domain important for protein-protein interaction. This protein was shown to	
	interact with mitotic checkpoint proteins including Mad2, p55CDC and BUBR1, and thus may be	
	involved in controlling the timing of mitosis. Alternatively spliced transcript variants encoding	
	different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]	
Gene ID:	996	
Application Details		
Application Notes:	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. ProClin	

Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months