

Datasheet for ABIN757584

anti-CARM1 antibody (pSer228) (Biotin)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	CARM1
Binding Specificity:	pSer228
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CARM1 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human CARM1 around the phosphorylation site of Ser228
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	CARM1
Alternative Name:	CARM1 (CARM1 Products)

Target Details

Background:	Synonyms: CARM1_HUMAN, Coactivator associated arginine methyltransferase 1, Coactivator-associated arginine methyltransferase 1, Histone arginine methyltransferase CARM 1, Histone arginine methyltransferase CARM1, Histone-arginine methyltransferase CARM1, PRMT 4, Protein arginine methyltransferase, Protein arginine N methyltransferase 4, Protein arginine N-methyltransferase 4. Background: Methylates (mono- and asymmetric dimethylation) the guanidino nitrogens of arginyl residues in several proteins involved in DNA packaging, transcription regulation, and mRNA stability. Recruited to promoters upon gene activation together with histone acetyltransferases from EP300/P300 and p160 families, methylates histone H3 at 'Arg-17' and activates transcription via chromatin remodeling. During nuclear hormone receptor activation and TCF7L2/TCF4 activation, acts synergically with EP300/P300 and either one of the p160 histone acetyltransferases NCOA1/SRC1, NCOA2/GRIP1 and NCOA3/ACTR or CTNNB1/beta-catenin to activate transcription. During myogenic transcriptional activation, acts together with NCOA3/ACTR as a coactivator for MEF2C.
Gene ID:	10498
UniProt:	Q86X55
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway , Regulation of Intracellular Steroid Hormone Receptor Signaling , Regulation of Lipid Metabolism by PPARalpha , Regulation of Muscle Cell Differentiation , Skeletal Muscle Fiber Development , Positive Regulation of fat Cell Differentiation

Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months