# ANTIBODIES ONLINE

## Datasheet for ABIN7581791 **Recombinant Mouse anti-Human IgG (Fc Region) Antibody** (HRP)





## Overview

000101000	
Quantity:	500 μL
Target:	IgG
Binding Specificity:	Fc Region
Reactivity:	Human
Host:	Mouse
Expression System:	Phage display
Antibody Type:	Recombinant Antibody
Clonality:	Multiclonal
Conjugate:	HRP
Application:	ELISA, Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (IF)
Product Details	
Purpose:	recombinant Multiclonal anti-human IgG (Fc-specific)
Immunogen:	No immunization, animal-free antibody development. Antigen: human intravenous immunoglobulins (IVIg)
lsotype:	lgG2a
Specificity:	This is an antibody developed by antibody phage display technology using a human naive antibody gene library and human IgG antigen. For this antibody both the heavy and light chains are cloned and expressed, generating full-length antibodies.
No Cross-Reactivity:	Cat, Cow, Dog, Goat, Horse, Human, Mouse, Rabbit, Rat

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7581791 | 07/11/2025 | Copyright antibodies-online. All rights reserved.

Product Details	
Cross-Reactivity (Details):	No known cross reactivity. No cross-reactivity to human IgM, human IgE, human IgA, mouse IgG, rabbit IgG, horse IgG, bovine IgG, goat IgG, rat IgG, dog IgG, cat IgG. Other species were not tested.
Characteristics:	Multiclonals are recombinant secondary antibodies that combine the best of polyclonal antisera and hybridoma monoclonal antibodies, while eliminating their disadvantages, plus they add the quality of recombinant reagents. Multiclonal antibodies consist of carefully adjusted mixtures recognizing different epitopes on all four different subclasses of human IgG. Their respective epitope binding sites do not compete with each other, therefore amplifying signal strengths. Multiclonals can contain up to 17 different individually tested monoclonal recombinant antibodies. This provides the typical advantage of polyepitope recognition which is key to the broad application profile of polyclonal antisera, but eliminates their disadvantages (limited batch sizes and batch-to-batch variations, no long-term reproducibility, undefined composition, unknown constituents). Multiclonals are completely sequence defined, implying their unlimited long-term availability and always identical test results. Their composition of individually characterised antibodies minimizes cross-reactivity with other targets, since they do not contain non-target directed IgG like all animal derived polyclonals do. This allows for a much lower unspecific binding reactivity in many assays compared to animal based products.
Purification:	Protein A purification
Grade:	Animal-Free

## Target Details

Target:	lgG
Abstract:	IgG Products
Target Type:	Antibody
Molecular Weight:	150 kDa
UniProt:	P01857

## Application Details

Restrictions:	For Research Use only
	determined by the investigator
Application Notes:	Western Blot: 1:1,000 - 1:20,000 ELISA: 1:1,000 - 1:50,000 Optimal working dilution should be

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7581791 | 07/11/2025 | Copyright antibodies-online. All rights reserved.

#### Handling

Format:	Liquid
Buffer:	PBS, pH 7.4, 0,02 % ProClin300
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C

#### Images



#### **ELISA**

**Image 1.** Different human IgG subclasses were coated on an ELISA plate at 100 ng. Multiclonal anti-human IgG (Fcspecific), (ABIN7581791), was titrated on top. Binding reactions were visualized using TMB. Absorbance was measured in an ELISA plate reader at 450 nm using the signal at 620 nm as reference.

anti-human IgG; Fc-specific



- Human IgG (polyclonal)
   IgY (monoclonal)
- bovine IgG (polyclonal)
- rabbit IgG (polyclonal)

   **▼** goat IgG (polyclonal)
- + horse IgG (polyclonal)
- ▲ rat IgG (polyclonal)

### ELISA

**Image 2.** Different species IgG were coated on an ELISA plate at 100 ng. Multiclonal anti-human IgG (Fc-specific), (ABIN7581791), was titrated on top. Binding reactions were visualized using TMB. Absorbance was measured in an ELISA plate reader at 450 nm using the signal at 620 nm as reference.