

## Datasheet for ABIN758192 anti-FGFR1 antibody (pTyr307) (PE)



( )	11 /	er	~\ /	10	1 A /
	1 / /	-1	١/	-	1///
$\sim$	٧.	$\sim$ 1	v	$\sim$	· v v

Overview	
Quantity:	100 μL
Target:	FGFR1
Binding Specificity:	pTyr307
Reactivity:	Human, Mouse, Rat, Cow, Chicken, Pig, Guinea Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FGFR1 antibody is conjugated to PE
Application:	Western Blotting (WB)
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human FGFR1 around the
	phosphorylation site of Tyr307
Isotype:	lgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Cow,Pig,Horse,Chicken,Guinea Pig
Purification:	Purified by Protein A.
Target Details	
Target:	FGFR1
Alternative Name:	FGFR1 (FGFR1 Products)

Background:
-------------

Synonyms: FGFR1 phospho Y307, p-FGFR1 phospho Y307, bFGFR, BFGFR, C FGR, CD 331, CD331, CD331 antigen, CEK, FGFBR, FGFR 1, Fibroblast growth factor receptor 1, FLG, FLG protein, FLJ14326, FLT 2, FLT2, Fms like tyrosine kinase 2, Fms related tyrosine kinase 2, Fms related tyrosine kinase 2 Pfefer syndrome, H2, H3, H4, H5, HBGFR, Heparin binding growth factor receptor, Hydroxyaryl protein kinase, KAL 2, KAL 2, MFR, N SAM, N sam tyrosine kinase, Protein tyrosine kinase, Tyrosylprotein kinase, Basic fibroblast growth factor receptor 1. Background: Fibroblast growth factors (FGFs) produce mitogenic and angiogenic effects in target cells by signaling through the cellular surface tyrosine kinase receptors. There are four members of the FGF receptor family: FGFR-1 (flg), FGFR-2 (bek, KGFR), FGFR-3 and FGFR-4. Each receptor contains an extracellular ligand binding domain, a transmembrane region and a cytoplasmic kinase domain (1). Following ligand binding and dimerization, the receptors are phosphorylated at specific tyrosine residues (2). Seven tyrosine residues in the cytoplasmic tail of FGFR-1 can be phosphorylated: Tyr463, Tyr583, Tyr585, Tyr653, Tyr654, Tyr730 and Tyr766. Tyrosine 653 and 654 are important for catalytic activity of the activated FGFR and are essential for signaling (3). The other phosphorylated tyrosine residues may provide docking sites for downstream signaling components such as Crk and PLCgamma.

Signaling Pathway, Sensory Perception of Sound, Stem Cell Maintenance, S100 Proteins

Molecular Weight:	90kDa
Gene ID:	2260
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin

## **Application Details**

Restrictions: For Research Use only

## Handling

Hariding	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months