

Datasheet for ABIN7582002

anti-ADORA1 antibody (Extracellular) (FITC)



[Go to Product page](#)

Overview

Quantity:	50 µL
Target:	ADORA1
Binding Specificity:	AA 153-168, Extracellular
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ADORA1 antibody is conjugated to FITC
Application:	Flow Cytometry (FACS), Live Cell Imaging (LCI)

Product Details

Purpose:	A Rabbit Polyclonal Antibody to Adenosine A1 Receptor (extracellular) conjugated to the fluorescent dye FITC
Immunogen:	(C)EQDWRANGSVGEPVIK, corresponding to amino acid residues 153 - 168 of rat Adenosine A1 Receptor
Sequence:	(C)EQDWRANGSV GEPVIK
Isotype:	IgG
Specificity:	Extracellular, 2nd loop.
Predicted Reactivity:	Mouse - 14 out of 16 amino acid residues identical Not recommended for human samples
Characteristics:	Anti-Adenosine A1 Receptor (extracellular) Antibody (ABIN7581822) is a highly specific antibody directed against an extracellular epitope of the rat protein. The antibody can be used in

Product Details

western blot, immunohistochemistry and flow cytometry applications. It has been designed to recognize Adenosine A1 Receptor from mouse and rat samples. The antibody is not recommended for human samples. Anti-Adenosine A1 Receptor (extracellular)-FITC Antibody (ABIN7581822)-F is directly conjugated to fluorescein isothiocyanate (FITC) fluorophore. This conjugated antibody has been developed to be used in immunofluorescent applications such as direct flow cytometry and live cell imaging.

Purification: Affinity purified on immobilized antigen.

Target Details

Target: ADORA1

Alternative Name: ADORA1 ([ADORA1 Products](#))

Background: Adenosine Receptor A1, Adenosine A1R, ADORA1, Adenosine is an endogenous nucleoside generated locally in tissues under conditions of hypoxia, ischemia, or inflammation. It modulates a variety of physiological functions in many tissues including the brain and heart.^{1,2} Adenosine exerts its actions via four specific adenosine receptors (also named P1 purinergic receptors): Adenosine A1 Receptor (A1AR), Adenosine A2A Receptor (A2AAR), Adenosine A2B Receptor (A2BAR), and Adenosine A3 Receptor (A3AR). All are integral membrane proteins and are members of the G protein-coupled receptor superfamily. They share a common structure of seven putative transmembrane domains, an extracellular amino terminus, a cytoplasmic carboxyl terminus, and a third intracellular loop that is important for binding G proteins.¹⁻³ The adenosine receptors can be distinguished on the basis of their differential selectivity for adenosine analogs.¹⁻³ A1AR is widely distributed and has been well characterized. High expression of A1AR is found in the brain (mainly in the cortex, cerebellum, and hippocampus), dorsal horn of the spinal cord, adrenal gland, and atria, and to a lower extent in several other tissues including adipose tissue, the colon, and kidney.^{2,4} A1AR modulates the activity of several ion channels. Activation of A1AR (by adenosine, its major agonist) inhibits N-type Ca²⁺ channels via a voltage-dependent, pertussis toxin (PTX)-sensitive pathway in neurons of the rat major pelvic ganglia (MPG).⁵ Since A1AR is the most prominent adenosine receptor in adipocytes, it has become a natural target for research on obesity, which is a major health problem.^{6,7} A possible role in cell proliferation and carcinogenesis has also been suggested for A1AR.^{8,9}

Gene ID: 29290

UniProt: [P25099](#)

Target Details

Pathways: [EGFR Signaling Pathway](#), [Negative Regulation of Hormone Secretion](#), [Synaptic Membrane](#)

Application Details

Application Notes: Antigen preadsorption control: 1 µg peptide per 1 µg antibody
Application Dilutions Immunohistochemistry paraffin embedded sections ihc: N/A
Application Dilutions Western blot wb: N/A

Comment: Negative Control: (ABIN7582044)
Blocking Peptide:

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: 50 µL double distilled water (DDW).

Concentration: 1 mg/mL

Buffer: PBS pH 7.4, 1 % BSA with 0.05 % sodium azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Storage before reconstitution: The antibody ships as a lyophilized powder at room temperature. Upon arrival, it should be stored at -20°C.
Storage after reconstitution: The reconstituted solution can be stored at 4°C, protected from the light, for up to 1 week. For longer periods, small aliquots should be stored at -20°C. Avoid multiple freezing and thawing. Centrifuge all antibody preparations before use (10000 x g 5 min).