

Datasheet for ABIN758481

anti-PCDHGB5 antibody (PE)



Overview

| Quantity: | 100 μL |
|--------------|-------------------------------------------|
| Target: | PCDHGB5 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PCDHGB5 antibody is conjugated to PE |
| Application: | Western Blotting (WB) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human PCDHGB5 |
|-----------------------|-------------------------------------------------------------|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat |
| Purification: | Purified by Protein A. |

Target Details

| Target: | PCDHGB5 |
|-------------------|---------------------------------------------------------------------------------------|
| Alternative Name: | Pcdhgb5 (PCDHGB5 Products) |
| Background: | Synonyms: PCDH gamma B5, PCDHGB 5, Protocadherin gamma B5, Protocadherin gamma |
| | subfamily B 5, PCDGH_HUMAN. |
| | Background: PCDHGB5 is a member of the protocadherin gamma gene cluster, one of three |
| | related clusters tandemly linked on chromosome five. These gene clusters have an |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn | International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com | Page 1/2 | Product datasheet for ABIN758481 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

immunoglobulin-like organization, suggesting that a novel mechanism may be involved in their regulation and expression. The gamma gene cluster includes 22 genes divided into 3 subfamilies. Subfamily A contains 12 genes, subfamily B contains 7 genes and 2 pseudogenes, and the more distantly related subfamily C contains 3 genes. The tandem array of 22 large, variable region exons are followed by a constant region, containing 3 exons shared by all genes in the cluster. Each variable region exon encodes the extracellular region, which includes 6 cadherin ectodomains and a transmembrane region. The constant region exons encode the common cytoplasmic region. These neural cadherin-like cell adhesion proteins most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been described for the gamma cluster genes.

Application Details

| Application Notes: | FCM(1:20-100) |
|--------------------|---------------|
| | |

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|-------------------------------------------------------------------------------------------------------------------------|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |