

Datasheet for ABIN7587670

PPIC Protein (AA 1-212) (His tag)



Overview

Quantity:	100 μg	
Target:	PPIC	
Protein Characteristics:	AA 1-212	
Origin:	Cow	
Source:	Yeast	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This PPIC protein is labelled with His tag.	
Application:	ELISA	
Product Details		
Sequence:	MGPGLRPLLP LVLCVGLSAL VPSAGASGFR KRGPSVTAKV FFDVRIGDKD VGRIVIGLFG	
	KVVPKTVENF VALATGEKGY GYKGSKFHRV IKDFMIQGGD FTRGDGTGGI SIYGETFPDE	
	NFKLKHYGIG WVSMANAGPD TNGSQFFITL TKPTWLDGKH VVFGKVLDGM TVVHSIELQA	
	TDGHDRPFTD CSIVNSGKID VKTPFVVEVS DW	
Specificity:	Bos taurus (Bovine)	
Characteristics:	Please inquire if you are interested in this recombinant protein expressed in E. coli, mammalien	
	cells or by baculovirus infection. Be aware about differences in price and lead time.	
Purity:	> 90 %	
Target Details		
Target:	PPIC	

Target Details

Alternative Name:	Peptidyl-prolyl cis-trans isomerase C (PPIC) (PPIC Products)
Background:	Recommended name: Peptidyl-prolyl cis-trans isomerase C.
	Short name= PPlase C.
	EC= 5.2.1.8.
	Alternative name(s): Cyclophilin C Rotamase C
UniProt:	Q08E11

Application Details

\sim				
Co	m	m	ıΔr	١т.
		11	1	II.

The yeast protein expression system is the most economical and efficient eukaryotic system for secretion and intracellular expression. A protein expressed by the mammalian cell system is of very high-quality and close to the natural protein. But the low expression level, the high cost of medium and the culture conditions restrict the promotion of mammalian cell expression systems. The yeast protein expression system serve as a eukaryotic system integrate the advantages of the mammalian cell expression system. A protein expressed by yeast system could be modificated such as glycosylation, acylation, phosphorylation and so on to ensure the native protein conformation. It can be used to produce protein material with high added value that is very close to the natural protein. Our proteins produced by yeast expression system has been used as raw materials for downstream preparation of monoclonal antibodies.

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Concentration:	0.2-2 mg/mL
Buffer:	Tris-based buffer, 50 % glycerol
Handling Advice:	Repeated freezing and thawing is not recommended. Store working aliquots at 4 °C for up to one week
Storage:	-20 °C
Storage Comment:	Store at -20 °C, for extended storage, conserve at -20 °C or -80 °C.