antibodies .- online.com







anti-MUC7 antibody (AA 21-120) (Cy5)



| () | ve | K\ / | | |
|-----|--------|------|----|----|
| | \cup | 1 V/ | Щ. | V۷ |
| | | | | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | MUC7 |
| Binding Specificity: | AA 21-120 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MUC7 antibody is conjugated to Cy5 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human MUC7 |
|-----------------------|--|
| Isotype: | IgG |
| Predicted Reactivity: | Human |
| Purification: | Purified by Protein A. |

Target Details

| Target: | MUC7 |
|--|----------------------|
| Alternative Name: | MUC7 (MUC7 Products) |
| Background: Synonyms: MG2, Mucin-7, MUC-7, Apo-MG2, Salivary mucin-7, MUC7 | |

Target Details

| l arget Details | |
|---------------------|--|
| | Background: May function in a protective capacity by promoting the clearance of bacteria in the |
| | oral cavity and aiding in mastication, speech, and swallowing. Binds P.aeruginosa pili. |
| Gene ID: | 4589 |
| UniProt: | Q8TAX7 |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |