

Datasheet for ABIN759508

anti-HIST1H3A antibody (AA 131-230) (Biotin)



Overview

Quantity:	100 μL
Target:	HIST1H3A
Binding Specificity:	AA 131-230
Reactivity:	Human, Mouse, Rat, Sheep
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HIST1H3A antibody is conjugated to Biotin
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HIF3 alpha
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat, Sheep
Predicted Reactivity:	Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	HIST1H3A
Alternative Name:	H3 alpha (HIST1H3A Products)

Target Details

Bac	kar	'nΙ	ın	Ч.
Duo		\sim	<i>.</i>	ч.

Synonyms: IPAS, MOP7, PASD7, HIF-3A, bHLHe17, Hypoxia-inducible factor 3-alpha, HIF-3-alpha, Basic-helix-loop-helix-PAS protein MOP7, Class E basic helix-loop-helix protein 17, HIF3-alpha-1, Inhibitory PAS domain protein, Member of PAS protein 7, PAS domain-containing protein 7, HIF3A

Background: Involved in adaptive response to hypoxia. Suppresses hypoxia-inducible expression of HIF1A and EPAS1. Binds to core DNA sequence 5'-TACGTG-3' within the hypoxia response element (HRE) of target gene promoters. The complex HIF3A-ARNT activates the transcription of reporter genes driven by HRE. Isoform 4 has a dominant-negative function of inactivating HIF1A-mediated transcription. Isoform 4 attenuates the binding of HIF1A to hypoxia-responsive elements (HRE), thus inhibiting HRE-driven transcription. Hypoxia induces down-regulation of isoform 4, leading to activation of HIF1A in hypoxia. Conversely, upon restoring normoxia, the expression of isoform 4 increases and thereby secure an inhibition of HIF1A activity. Isoform 4 may be a negative regulator of hypoxia-inducible gene expression in the kidney and may be involved in renal tumorigenesis. Functions as an inhibitor of angiogenesis in the cornea (By similarity).

Gene ID:

64344

UniProt:

Q9Y2N7

Application Details

Application Notes:

WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

Handling

_	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months