

Datasheet for ABIN7596771

GPR18 Protein (DYKDDDDK Tag, Strep Tag)



[Go to Product page](#)

Overview

Quantity:	10 µg
Target:	GPR18
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Synthetic Nanodisc
Purification tag / Conjugate:	This GPR18 protein is labelled with DYKDDDDK Tag, Strep Tag.
Application:	ELISA, Surface Plasmon Resonance (SPR), Phage Display (PhD), Immunogen (Imm), Cryogenic electron microscopy (cryo-EM)

Product Details

Purpose:	Human GPR18-Strep full length protein-synthetic nanodisc
----------	--

Target Details

Target:	GPR18
Alternative Name:	GPR18 (GPR18 Products)
Background:	N-arachidonyl glycine receptor, NAGly receptor, G-protein coupled receptor 18, GPCR18, GPR18 Receptor for endocannabinoid N-arachidonyl glycine (NAGly) (PubMed:16844083, PubMed:24762058, PubMed:27572937). However, conflicting results about the role of NAGly as an agonist are reported (PubMed:27018161). Can also be activated by plant-derived and synthetic cannabinoid agonists (PubMed:24762058). The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase (PubMed:16844083). May contribute to regulation of the immune system. Is required for normal homeostasis of CD8 subsets of intraepithelial

Target Details

lymphocytes (IELs) (CD8alphaalpha and CD8alphabeta IELs)in small intestine by supporting preferential migration of CD8alphaalpha T-cells to intraepithelial compartment over lamina propria compartment, and by mediating their reconstitution into small intestine after bone marrow transplant (By similarity). Plays a role in hypotensive responses, mediating reduction in intraocular and blood pressure (By similarity). Mediates NAGly-induced process of reorganization of actin filaments and induction of acrosomal exocytosis (PubMed:27572937).[UniProtKB/Swiss-Prot Function]

Molecular Weight: The human full length GPR18-Strep protein has a MW of 38.1 kDa

UniProt: [Q14330](#)

Application Details

Comment: Advantages:

- Highly purified membrane proteins
- High solubility in aqueous solutions
- High stability
- Proteins are in a native membrane environment and remain biologically active
- No detergent and can be used for cell-based assays
- No MSP backbone proteins
- Mammalian cell expression system ensures post- translational modifications

Restrictions: For Research Use only

Handling

Format: Lyophilized

Buffer: Solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

Storage: -20 °C,-80 °C

Storage Comment: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

Expiry Date: 12 months