

Datasheet for ABIN7597083

KCTD11 Protein (DYKDDDDK Tag,Strep Tag)



Go to Product page

_			
()	V/C	rv	٨/

Quantity:	10 μg	
Target:	KCTD11	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Synthetic Nanodisc	
Purification tag / Conjugate:	This KCTD11 protein is labelled with DYKDDDDK Tag,Strep Tag.	
Application:	ELISA, Cryogenic electron microscopy (cryo-EM), Immunogen (Imm), Phage Display (PhD), Surface Plasmon Resonance (SPR)	
Product Details		
Purpose:	Human KCD11-Strep full length protein-synthetic nanodisc	
Target Details		
Target:	KCTD11	
Alternative Name:	KCD11 (KCTD11 Products)	
Background:	C17orf36, KCASH1, REN, REN/KCTD11	
	Plays a role as a marker and a regulator of neuronal differentiation, Up-regulated by a variety of	
	neurogenic signals, such as retinoic acid, epidermal growth factor/EGF and NGFB/nerve growth	
	factor. Induces apoptosis, growth arrest and the expression of cyclin-dependent kinase inhibitor	
	CDKN1B. Plays a role as a tumor repressor and inhibits cell growth and tumorigenicity of	
	medulloblastoma (MDB). Acts as probable substrate-specific adapter for a BCR (BTB-CUL3-	
	RBX1) E3 ubiquitin-protein ligase complex towards HDAC1. Functions as antagonist of the	

Target Details

	Hedgehog pathway on cell proliferation and differentiation by affecting the nuclear transf	
	transcription factor GLI1, thus maintaining cerebellar granule cells in undifferentiated state, this	
	effect probably occurs via HDAC1 down-regulation, keeping GLI1 acetylated and inactive. When	
	knock-down, Hedgehog antagonism is impaired and proliferation of granule cells is sustained.	
	Activates the caspase cascade.[UniProtKB/Swiss-Prot Function]	
Molecular Weight:	The human full length KCD11-Strep protein has a MW of 25.9 kDa	
UniProt:	Q693B1	
Pathways:	Hedgehog Signaling	

Application Details

\sim			m+.
Co	1111	116	. 111

Advantages:

- · Highly purified membrane proteins
- · High solubility in aqueous solutions
- High stability
- · Proteins are in a native membrane environment and remain biologically active
- · No detergent and can be used for cell-based assays
- No MSP backbone proteins
- · Mammalian cell expression system ensures post-translational modifications

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Buffer:	Solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Expiry Date:	12 months