

Datasheet for ABIN7597319

E-cadherin Protein (AA 263-375) (Fc Tag)



Go to Product page

()	ve	r\/i	Δ	۱۸/
\circ	V C	1 V		v v

Quantity:	10 μg
Target:	E-cadherin (CDH1)
Protein Characteristics:	AA 263-375
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This E-cadherin protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant human CDH1(263-375) Protein with C-terminal human Fc tag	
Sequence:	CDH1(Thr263-Phe375) hFc(Glu99-Ala330)	
Purity:	The purity of the protein is greater than 90 % as determined by SDS-PAGE and Coomassie blue	
	staining.	

Target Details

Target:	E-cadherin (CDH1)	
Alternative Name:	CDH1 (CDH1 Products)	
Background:	UVO, CDHE, ECAD, LCAM, Arc-1, BCDS1, CD324	
	This gene encodes a classical cadherin of the cadherin superfamily. Alternative splicing results	
	in multiple transcript variants, at least one of which encodes a preproprotein that is	
	proteolytically processed to generate the mature glycoprotein. This calcium-dependent cell-cell	

	adhesion protein is comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Mutations in this gene are correlated with gastric, breast, colorectal, thyroid and ovarian cancer. Loss of function of this gene is thought to contribute to cancer progression by increasing proliferation, invasion, and/or metastasis. The ectodomain of this protein mediates bacterial adhesion to mammalian cells and the cytoplasmic domain is required for internalization. This gene is present in a gene cluster with other members of the cadherin family on chromosome 16. [provided by RefSeq, Nov 2015]	
Molecular Weight:	predicted molecular mass of 38.1 kDa after removal of the signal peptide.	
UniProt:	P12830	
Pathways:	WNT Signaling, Sensory Perception of Sound, Cell-Cell Junction Organization, Tube Formation	
Application Details		
Application Notes:	 Extracellular Domain Proteins (ECD) can be used as: Immunogens for antibody drug development Reagents used for CAR-T positive cell monitoring Reagents for antibody screening and functional testing Reagents for antibody affinity measurement 	
Comment:	The protein was made using HEK293 mammalian cell secretion expression system to ensure the close-to-native structures and post-translational modifications of the target protein.	
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Buffer:	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.	
Expiry Date:	12 months	