

Datasheet for ABIN7597332

## c-MET Protein (AA 740-932) (Fc Tag)



[Go to Product page](#)

### Overview

Quantity:	10 µg
Target:	c-MET (MET)
Protein Characteristics:	AA 740-932
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This c-MET protein is labelled with Fc Tag.

### Product Details

Purpose:	Recombinant human MET(740-932) Protein with C-terminal human Fc tag
Sequence:	MET(Glu740-Thr932) hFc(Glu99-Ala330)
Purity:	The purity of the protein is greater than 95 % as determined by SDS-PAGE and Coomassie blue staining.

### Target Details

Target:	c-MET (MET)
Alternative Name:	MET ( <a href="#">MET Products</a> )
Background:	<p>DA11, HGFR, AUTS9, RCCP2, c-Met, DFNB97</p> <p>This gene encodes a member of the receptor tyrosine kinase family of proteins and the product of the proto-oncogene MET. The encoded preproprotein is proteolytically processed to generate alpha and beta subunits that are linked via disulfide bonds to form the mature receptor. Further</p>

## Target Details

processing of the beta subunit results in the formation of the M10 peptide, which has been shown to reduce lung fibrosis. Binding of its ligand, hepatocyte growth factor, induces dimerization and activation of the receptor, which plays a role in cellular survival, embryogenesis, and cellular migration and invasion. Mutations in this gene are associated with papillary renal cell carcinoma, hepatocellular carcinoma, and various head and neck cancers. Amplification and overexpression of this gene are also associated with multiple human cancers. [provided by RefSeq, May 2016]

**Molecular Weight:** predicted molecular mass of 47.5 kDa after removal of the signal peptide. The apparent molecular mass of MET(740-932)-hFc is 55-70 kDa due to glycosylation.

**UniProt:** [P08581](#)

**Pathways:** [RTK Signaling](#), [Carbohydrate Homeostasis](#), [Synaptic Membrane](#), [Signaling of Hepatocyte Growth Factor Receptor](#)

## Application Details

**Application Notes:** Extracellular Domain Proteins (ECD) can be used as:

- Immunogens for antibody drug development
- Reagents used for CAR-T positive cell monitoring
- Reagents for antibody screening and functional testing
- Reagents for antibody affinity measurement

**Comment:** The protein was made using HEK293 mammalian cell secretion expression system to ensure the close-to-native structures and post-translational modifications of the target protein.

**Restrictions:** For Research Use only

## Handling

**Format:** Lyophilized

**Buffer:** Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization.

**Storage:** -20 °C, -80 °C

**Storage Comment:** Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

**Expiry Date:** 12 months