

## Datasheet for ABIN7597365 EGFR Protein (AA 25-389) (His tag)



_				
( )	1//	rv	IO	Λ/
( )	VC	. I V	1	v v

Quantity:	10 μg
Target:	EGFR
Protein Characteristics:	AA 25-389
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This EGFR protein is labelled with His tag.

## **Product Details**

Purpose:	Recombinant human EGFR(25-389) Protein with C-terminal 10xHis tag	
Sequence:	EGFR(Leu25-Pro389) 10xHis tag	
Purity:	The purity of the protein is greater than 85 % as determined by SDS-PAGE and Coomassie blu	
	staining.	

## **Target Details**

Target:	EGFR	
Alternative Name:	EGFR (EGFR Products)	
Background:	ERBB, ERRP, HER1, mENA, ERBB1, NNCIS, PIG61, NISBD2	
	The protein encoded by this gene is a transmembrane glycoprotein that is a member of the	
	protein kinase superfamily. This protein is a receptor for members of the epidermal growth	
	factor family. EGFR is a cell surface protein that binds to epidermal growth factor, thus inducing	

	receptor dimerization and tyrosine autophosphorylation leading to cell proliferation. Mutations	
	in this gene are associated with lung cancer. EGFR is a component of the cytokine storm which	
	contributes to a severe form of Coronavirus Disease 2019 (COVID-19) resulting from infection	
	with severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2). [provided by RefSeq, Jul	
	2020]	
Molecular Weight:	predicted molecular mass of 42.0 kDa after removal of the signal peptide. The apparent	
	molecular mass of EGFR(25-389)-His is 35-70 kDa due to glycosylation.	
UniProt:	P00533	
Pathways:	NF-kappaB Signaling, RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling	
	Pathway, Neurotrophin Signaling Pathway, Stem Cell Maintenance, Hepatitis C, Positive	
	Regulation of Response to DNA Damage Stimulus, Interaction of EGFR with phospholipase C-	
	gamma, Thromboxane A2 Receptor Signaling, EGFR Downregulation, S100 Proteins	
Application Details		
Application Notes:	Extracellular Domain Proteins (ECD) can be used as:	
	Immunogens for antibody drug development	
	Reagents used for CAR-T positive cell monitoring	
	Reagents for antibody screening and functional testing	
	Reagents for antibody affinity measurement	
Comment:	The protein was made using HEK293 mammalian cell secretion expression system to ensure	
	the close-to-native structures and post-translational modifications of the target protein.	
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Buffer:	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants	
	before lyophilization.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for	
	use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).	
	Lyophilized proteins are shipped at ambient temperature.	
Expiry Date:	12 months	