

Datasheet for ABIN7597414

ADAM17 Protein (AA 18-671) (Fc Tag)



[Go to Product page](#)

Overview

Quantity:	10 µg
Target:	ADAM17
Protein Characteristics:	AA 18-671
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ADAM17 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant human ADAM17 Protein with C-terminal human Fc tag
Sequence:	ADAM17(Pro18-Asn671) hFc(Glu99-Ala330)
Purity:	The purity of the protein is greater than 95 % as determined by SDS-PAGE and Coomassie blue staining.

Target Details

Target:	ADAM17
Alternative Name:	ADAM17 (ADAM17 Products)
Background:	<p>CSVP, TACE, CD156B, NISBD1</p> <p>This gene encodes a member of the ADAM (a disintegrin and metalloprotease domain) family. Members of this family are membrane-anchored proteins structurally related to snake venom disintegrins, and have been implicated in a variety of biologic processes involving cell-cell and</p>

Target Details

cell-matrix interactions, including fertilization, muscle development, and neurogenesis. The encoded preproprotein is proteolytically processed to generate the mature protease. The encoded protease functions in the ectodomain shedding of tumor necrosis factor-alpha, in which soluble tumor necrosis factor-alpha is released from the membrane-bound precursor. This protease also functions in the processing of numerous other substrates, including cell adhesion proteins, cytokine and growth factor receptors and epidermal growth factor (EGF) receptor ligands, and plays a prominent role in the activation of the Notch signaling pathway. Elevated expression of this gene has been observed in specific cell types derived from psoriasis, rheumatoid arthritis, multiple sclerosis and Crohn's disease patients, suggesting that the encoded protein may play a role in autoimmune disease. Additionally, this protease may play a role in viral infection through its cleavage of ACE2, the cellular receptor for SARS-CoV and SARS-CoV-2. [provided by RefSeq, Aug 2020]

Molecular Weight:	predicted molecular mass of 100.2 kDa after removal of the signal peptide.
UniProt:	P78536
Pathways:	Notch Signaling , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Response to Growth Hormone Stimulus

Application Details

Application Notes:	Extracellular Domain Proteins (ECD) can be used as: <ul style="list-style-type: none">• Immunogens for antibody drug development• Reagents used for CAR-T positive cell monitoring• Reagents for antibody screening and functional testing• Reagents for antibody affinity measurement
Comment:	The protein was made using HEK293 mammalian cell secretion expression system to ensure the close-to-native structures and post-translational modifications of the target protein.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Buffer:	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization.
Storage:	-20 °C, -80 °C

Handling

Storage Comment:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Expiry Date:	12 months
