

Datasheet for ABIN7597426

## B4GALT1 Protein (AA 45-398) (His tag)



[Go to Product page](#)

### Overview

Quantity:	10 µg
Target:	B4GALT1
Protein Characteristics:	AA 45-398
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This B4GALT1 protein is labelled with His tag.

### Product Details

Purpose:	Recombinant human B4GALT1 Protein with N-terminal 10xHis tag
Sequence:	10xHis tag B4GALT1(Arg45-Ser398)
Purity:	The purity of the protein is greater than 85 % as determined by SDS-PAGE and Coomassie blue staining.

### Target Details

Target:	B4GALT1
Alternative Name:	B4GALT1 ( <a href="#">B4GALT1 Products</a> )
Background:	<p>GT1, GTB, CDG2D, GGTB2, CLDLFIB, B4GAL-T1, beta4Gal-T1</p> <p>This gene is one of seven beta-1,4-galactosyltransferase (beta4GalT) genes. They encode type II membrane-bound glycoproteins that appear to have exclusive specificity for the donor substrate UDP-galactose, all transfer galactose in a beta1,4 linkage to similar acceptor sugars:</p>

## Target Details

GlcNAc, Glc, and Xyl. Each beta4GalT has a distinct function in the biosynthesis of different glycoconjugates and saccharide structures. As type II membrane proteins, they have an N-terminal hydrophobic signal sequence that directs the protein to the Golgi apparatus and which then remains uncleaved to function as a transmembrane anchor. By sequence similarity, the beta4GalTs form four groups: beta4GalT1 and beta4GalT2, beta4GalT3 and beta4GalT4, beta4GalT5 and beta4GalT6, and beta4GalT7. This gene is unique among the beta4GalT genes because it encodes an enzyme that participates both in glycoconjugate and lactose biosynthesis. For the first activity, the enzyme adds galactose to N-acetylglucosamine residues that are either monosaccharides or the nonreducing ends of glycoprotein carbohydrate chains. The second activity is restricted to lactating mammary tissues where the enzyme forms a heterodimer with alpha-lactalbumin to catalyze UDP-galactose + D-glucose UDP + lactose. The two enzymatic forms result from alternate transcription initiation sites and post-translational processing. Two transcripts, which differ only at the 5' end, with approximate lengths of 4.1 kb and 3.9 kb encode the same protein. The longer transcript encodes the type II membrane-bound, trans-Golgi resident protein involved in glycoconjugate biosynthesis. The shorter transcript encodes a protein which is cleaved to form the soluble lactose synthase. [provided by RefSeq, Jul 2008]

Molecular Weight: predicted molecular mass of 40.7 kDa after removal of the signal peptide.

UniProt: [P15291](#)

Pathways: [Glycosaminoglycan Metabolic Process](#)

## Application Details

Application Notes: Extracellular Domain Proteins (ECD) can be used as:

- Immunogens for antibody drug development
- Reagents used for CAR-T positive cell monitoring
- Reagents for antibody screening and functional testing
- Reagents for antibody affinity measurement

Comment: The protein was made using HEK293 mammalian cell secretion expression system to ensure the close-to-native structures and post-translational modifications of the target protein.

Restrictions: For Research Use only

## Handling

Format: Lyophilized

## Handling

---

Buffer:	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Expiry Date:	12 months