

Datasheet for ABIN7597469

MUC1 Protein (AA 380-500) (Fc Tag)



[Go to Product page](#)

Overview

Quantity:	10 µg
Target:	MUC1
Protein Characteristics:	AA 380-500
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MUC1 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Cynomolgus MUC1(380-500) protein with C-terminal human Fc tag
Sequence:	MUC1(Leu380-Val500) hFc(Glu99-Ala330)
Purity:	The purity of the protein is greater than 95 % as determined by SDS-PAGE and Coomassie blue staining.

Target Details

Target:	MUC1
Alternative Name:	MUC1 (MUC1 Products)
Background:	ADMCKD, ADMCKD1, CA 15-3, CD227, EMA, H23AG, KL-6, MAM6, MCD, MCKD, MCKD1, MUC-1, MUC-1/SEC, MUC-1/X, MUC1/ZD, PEM, PEMT, PUM This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on

Target Details

epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast stomach and pancreas. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results in multiple transcript variants.

Molecular Weight: predicted molecular mass of 39.8 kDa after removal of the signal peptide. The apparent molecular mass of cMUC1(380-500)-hFc is 35-55 kDa due to glycosylation.

NCBI Accession: [XP_005541632](#)

Pathways: [Negative Regulation of intrinsic apoptotic Signaling](#)

Application Details

Application Notes: Extracellular Domain Proteins (ECD) can be used as:

- Immunogens for antibody drug development
- Reagents used for CAR-T positive cell monitoring
- Reagents for antibody screening and functional testing
- Reagents for antibody affinity measurement

Comment: The protein was made using HEK293 mammalian cell secretion expression system to ensure the close-to-native structures and post-translational modifications of the target protein.

Restrictions: For Research Use only

Handling

Format: Lyophilized

Buffer: Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization.

Storage: -20 °C, -80 °C

Storage Comment: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).
Lyophilized proteins are shipped at ambient temperature.

Handling

Expiry Date: 12 months