

Datasheet for ABIN7598795

anti-Prostate Specific Antigen antibody (N-Term)

2 Images



Go to Product page

\sim				
O_1	/ el	rVI	161	Λ

Quantity:	7 mL
Target:	Prostate Specific Antigen (PSA)
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This Prostate Specific Antigen antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Purpose:	Anti DOA (Dunatata annaifin antinan)
i uipose.	Anti-PSA (Prostate-specific antigen)
Immunogen:	Peptide derived from N-terminal sequence of human PSA. Antibody recognizes the epitope between Arg77 -Gln88.
·	Peptide derived from N-terminal sequence of human PSA. Antibody recognizes the epitope
Immunogen:	Peptide derived from N-terminal sequence of human PSA. Antibody recognizes the epitope between Arg77 -Gln88.
Immunogen: Clone:	Peptide derived from N-terminal sequence of human PSA. Antibody recognizes the epitope between Arg77 -Gln88. L15-V
Immunogen: Clone: Isotype:	Peptide derived from N-terminal sequence of human PSA. Antibody recognizes the epitope between Arg77 -Gln88. L15-V IgG
Immunogen: Clone: Isotype: Specificity:	Peptide derived from N-terminal sequence of human PSA. Antibody recognizes the epitope between Arg77 -Gln88. L15-V IgG
Immunogen: Clone: Isotype: Specificity: Target Details	Peptide derived from N-terminal sequence of human PSA. Antibody recognizes the epitope between Arg77 -Gln88. L15-V IgG PSA (Prostate-specific antigen)

Target Details

Background:	PSA represents a prostate cancer protein marker, strongly expressed in both normal and
	neoplastic prostatic tissue. This protein is expressed at lower levels also in in breast tumors,
	endometrium, or renal cell carcinomas. Together with Prostate-Specific Acid Phosphatase
	(PSAP), PSA expression is detected in blood samples and/or prostate tissue-related biopsies, in
	order to confirm the prostatic origin of metastatic carcinoma.

UniProt: P07288

Pathways: Complement System


Application Details

Application Notes:	Ready to use antibody dilution. Directly use for application
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	20 mM Tris-HCl, pH 8.0, Stabilizer: 20 mg/mL BSA, Preservative: 0.05 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	+4°C, Do not freeze!

Images



Immunohistochemistry

Image 1.



Immunohistochemistry

Image 2.