

Datasheet for ABIN7598966

anti-HES5 antibody (AA 1-121)



\sim				
O_1	/ el	rVI	161	Λ

Quantity:	100 μg
Target:	HES5
Binding Specificity:	AA 1-121
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HES5 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Flow Cytometry (FACS)
Product Details	
Purpose:	Anti-HES5 Antibody Picoband®
Immunogen:	E.coli-derived human HES5 recombinant protein (Position: M1-R121).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-HES5 Antibody Picoband® (ABIN7598966). Tested in ELISA, Flow Cytometry, IHC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	HES5
Alternative Name:	HES5 (HES5 Products)
Background:	Synonyms: Cytoskeleton-associated protein 5, Colonic and hepatic tumor overexpressed gene
	protein, Ch-TOG, CKAP5, KIAA0097
	Tissue Specificity: Expressed in fetal brain. Highly expressed in brain and placenta. Lower levels
	in heart, liver, thymus, kidney and lung. Located to endothelial cells and neuronal cells of the
	suprachiasmatic nucleus (SCN). Also detected in endothelial cells of the heart, lung and kidney.
	In the brain, specifically expressed in the thalamus, hippocampus and amygdala.
	Background: Transcription factor HES-5 is a protein that in humans is encoded by the HES5
	gene. This gene encodes a member of a family of basic helix-loop-helix transcriptional
	repressors. The protein product of this gene, which is activated downstream of the Notch
	pathway, regulates cell differentiation in multiple tissues. Disruptions in the normal expression
	of this gene have been associated with developmental diseases and cancer.
Molecular Weight:	18 kDa
Gene ID:	388585
Pathways:	Stem Cell Maintenance, Tube Formation
·	Stem Cell Maintenance, Tube Formation
Application Details	
·	Western blot, 0.25-0.5 μg/mL, Human
Application Details	Western blot, 0.25-0.5 μg/mL, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 μg/mL, Human
Application Details	Western blot, 0.25-0.5 μg/mL, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 μg/mL, Human Flow Cytometry (Fixed), 1-3 μg/1x10 ⁶ cells, Human
Application Details	Western blot, 0.25-0.5 μg/mL, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 μg/mL, Human Flow Cytometry (Fixed), 1-3 μg/1x10 ⁶ cells, Human ELISA, 0.1-0.5 μg/mL, -
Application Details	Western blot, 0.25-0.5 μg/mL, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 μg/mL, Human Flow Cytometry (Fixed), 1-3 μg/1x10 ⁶ cells, Human ELISA, 0.1-0.5 μg/mL, - 1. Katoh, M., Katoh, M. Identification and characterization of human HES2, HES3, and HES5
Application Details	Western blot, 0.25-0.5 μg/mL, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 μg/mL, Human Flow Cytometry (Fixed), 1-3 μg/1x10 ⁶ cells, Human ELISA, 0.1-0.5 μg/mL, - 1. Katoh, M., Katoh, M. Identification and characterization of human HES2, HES3, and HES5 genes in silico. Int. J. Oncol. 25: 529-534, 2004. 2. Takebayashi, K., Akazawa, C., Nakanishi, S.,
Application Details	Western blot, 0.25-0.5 μg/mL, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 μg/mL, Human Flow Cytometry (Fixed), 1-3 μg/1x10 ⁶ cells, Human ELISA, 0.1-0.5 μg/mL, - 1. Katoh, M., Katoh, M. Identification and characterization of human HES2, HES3, and HES5 genes in silico. Int. J. Oncol. 25: 529-534, 2004. 2. Takebayashi, K., Akazawa, C., Nakanishi, S., Kageyama, R. Structure and promoter analysis of the gene encoding the mouse helix-loop-helix
Application Details	Western blot, 0.25-0.5 μg/mL, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 μg/mL, Human Flow Cytometry (Fixed), 1-3 μg/1x10 ⁶ cells, Human ELISA, 0.1-0.5 μg/mL, - 1. Katoh, M., Katoh, M. Identification and characterization of human HES2, HES3, and HES5 genes in silico. Int. J. Oncol. 25: 529-534, 2004. 2. Takebayashi, K., Akazawa, C., Nakanishi, S.,
Application Details	Western blot, 0.25-0.5 μg/mL, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 μg/mL, Human Flow Cytometry (Fixed), 1-3 μg/1x10 ⁶ cells, Human ELISA, 0.1-0.5 μg/mL, - 1. Katoh, M., Katoh, M. Identification and characterization of human HES2, HES3, and HES5 genes in silico. Int. J. Oncol. 25: 529-534, 2004. 2. Takebayashi, K., Akazawa, C., Nakanishi, S., Kageyama, R. Structure and promoter analysis of the gene encoding the mouse helix-loop-helix factor HES-5: identification of the neural precursor cell-specific promoter element. J. Biol.
Application Details Application Notes:	Western blot, 0.25-0.5 μg/mL, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 μg/mL, Human Flow Cytometry (Fixed), 1-3 μg/1x10 ⁶ cells, Human ELISA, 0.1-0.5 μg/mL, - 1. Katoh, M., Katoh, M. Identification and characterization of human HES2, HES3, and HES5 genes in silico. Int. J. Oncol. 25: 529-534, 2004. 2. Takebayashi, K., Akazawa, C., Nakanishi, S., Kageyama, R. Structure and promoter analysis of the gene encoding the mouse helix-loop-helix factor HES-5: identification of the neural precursor cell-specific promoter element. J. Biol. Chem. 270: 1342-1349, 1995.
Application Details Application Notes: Restrictions:	Western blot, 0.25-0.5 μg/mL, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 μg/mL, Human Flow Cytometry (Fixed), 1-3 μg/1x10 ⁶ cells, Human ELISA, 0.1-0.5 μg/mL, - 1. Katoh, M., Katoh, M. Identification and characterization of human HES2, HES3, and HES5 genes in silico. Int. J. Oncol. 25: 529-534, 2004. 2. Takebayashi, K., Akazawa, C., Nakanishi, S., Kageyama, R. Structure and promoter analysis of the gene encoding the mouse helix-loop-helix factor HES-5: identification of the neural precursor cell-specific promoter element. J. Biol. Chem. 270: 1342-1349, 1995.

Handling

Concentration:	500 μg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.