

Datasheet for ABIN7599404
anti-MEK2 antibody (AA 1-45)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	MEK2 (MAP2K2)
Binding Specificity:	AA 1-45
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MEK2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS)

Product Details

Purpose:	Anti-MEK2/MAP2K2 Antibody Picoband®
Immunogen:	E.coli-derived human MEK2/MAP2K2 recombinant protein (Position: M1-E45).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-MEK2/MAP2K2 Antibody Picoband® (ABIN7599404). Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	MEK2 (MAP2K2)
Alternative Name:	MAP2K2 (MAP2K2 Products)
Background:	Synonyms: 5'-AMP-activated protein kinase catalytic subunit alpha-1, AMPK subunit alpha-1 Tissue Specificity: Expressed in neurons of the rolandic area of the brain (at protein level). Expressed in the brain. Background: Dual specificity mitogen-activated protein kinase kinase 2(MAP2K2), also called PRKMK2 or MEK2, is an enzyme that in humans is encoded by the MAP2K2 gene. The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. MAP2K2 is mapped to 19p13.3. This kinase is known to play a critical role in mitogen growth factor signal transduction, and the inhibition or degradation of this kinase is found to be involved in the pathogenesis of Yersinia and anthrax. Recombinant MEK2 and MEK1 both could activate human ERK1 in vitro, and they further characterized biochemically the 2 MAP2Ks. MAP2K2 has been shown to interact with MAPK3 and ARAF.
Molecular Weight:	44 kDa
Gene ID:	5605
UniProt:	P36507
Pathways:	MAPK Signaling , RTK Signaling , Fc-epsilon Receptor Signaling Pathway , Neurotrophin Signaling Pathway , Activation of Innate immune Response , Toll-Like Receptors Cascades , Signaling of Hepatocyte Growth Factor Receptor , BCR Signaling

Application Details

Application Notes:	Western blot, 0.25-0.5 µg/mL, Human, Mouse, Rat Flow Cytometry (Fixed), 1-3 µg/1×10⁶ cells, Rat ELISA, 0.1-0.5 µg/mL, - 1."Entrez Gene: MAP2K2 mitogen-activated protein kinase kinase 2" 2. Zheng, C. F., Guan, K. L. Cloning and characterization of two distinct human extracellular signal-regulated kinase activator kinases, MEK1 and MEK2. J. Biol. Chem. 268: 11435-11439, 1993.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.

Handling

Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage:	4 °C, -20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.