

Datasheet for ABIN7599425

anti-SYT16 antibody (AA 1-474)



Overview

Quantity:	100 μg
Target:	SYT16
Binding Specificity:	AA 1-474
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SYT16 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF), Flow Cytometry (FACS)

Product Details

Purpose:	Anti-SYT16 Antibody Picoband®
Immunogen:	E.coli-derived human SYT16 recombinant protein (Position: M1-R474).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins
Characteristics:	Anti-SYT16 Antibody Picoband® (ABIN7599425). Tested in ELISA, Flow Cytometry, IF, ICC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	SYT16
Alternative Name:	SYT16 (SYT16 Products)
Background:	Synonyms: RNA-binding protein 47,RNA-binding motif protein 47,RBM47,
	Tissue Specificity: Abundantly expressed in tonsil, lymph node, and trachea, strong expression
	in prostate, lower expression in thyroid, stomach, and colon
	Background: Predicted to enable identical protein binding activity and phospholipid binding
	activity. Predicted to be involved in exocytosis. Predicted to be integral component of
	membrane.
Molecular Weight:	72 kDa
Gene ID:	83851

Application Details

Application Notes:	Western blot, 0.25-0.5 μg/mL, Human
	Immunocytochemistry/Immunofluorescence, 5 μg/mL, Human
	Flow Cytometry (Fixed), 1-3 µg/1x10 ⁶ cells, Human
	ELISA, 0.1-0.5 μg/mL, -
	1. Fukuda, M. Molecular cloning, expression, and characterization of a novel class of
	synaptotagmin (Syt XIV) conserved from Drosophila to humans. J. Biochem. 133: 641-649,
	2003.

For Research Use only

Handling

Restrictions:

Format:	Lyophilized
Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 μg/mL.
Concentration:	500 μg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.