

Datasheet for ABIN7599570
anti-DYNLL1 antibody (AA 1-89)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	DYNLL1
Binding Specificity:	AA 1-89
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This DYNLL1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Flow Cytometry (FACS), Immunocytochemistry (ICC)

Product Details

Purpose:	Anti-DYNLL1/PIN Antibody Picoband® (monoclonal, 6G2H1)
Immunogen:	E.coli-derived human DYNLL1/PIN recombinant protein (Position: M1-G89).
Clone:	6G2H1
Isotype:	IgG2b
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-DYNLL1/PIN Antibody Picoband® (monoclonal, 6G2H1) (ABIN7599570). Tested in Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.

Product Details

Purification: Immunogen affinity purified.

Target Details

Target: DYNLL1

Alternative Name: DYNLL1 ([DYNLL1 Products](#))

Background: Synonyms: Survival motor neuron protein, Component of gems 1, Gemin-1, SMN1, SMN, SMNT, SMN2, SMNC

Tissue Specificity: Expressed in a wide variety of tissues. Expressed at high levels in brain, kidney and liver, moderate levels in skeletal and cardiac muscle, and low levels in fibroblasts and lymphocytes. Also seen at high levels in spinal cord. Present in osteoclasts and mononuclear cells (at protein level).

Background: Dynein light chain 1, cytoplasmic is a protein that in humans is encoded by the DYNLL1 gene. Cytoplasmic dyneins are large enzyme complexes with a molecular mass of about 1,200 kD. They contain two force-producing heads formed primarily from dynein heavy chains, and stalks linking the heads to a basal domain, which contains a varying number of accessory intermediate chains. The complex is involved in intracellular transport and motility. The protein described in this record is a light chain and exists as part of this complex but also physically interacts with and inhibits the activity of neuronal nitric oxide synthase. Binding of this protein destabilizes the neuronal nitric oxide synthase dimer, a conformation necessary for activity, and it may regulate numerous biologic processes through its effects on nitric oxide synthase activity. Alternate transcriptional splice variants have been characterized.

Molecular Weight: 12 kDa

Gene ID: 8655

UniProt: [P63167](#)

Pathways: [M Phase](#), [Tube Formation](#), [Positive Regulation of Endopeptidase Activity](#)

Application Details

Application Notes: Western blot, 0.25-0.5 µg/mL, Human, Mouse, Rat
Immunohistochemistry(Paraffin-embedded Section), 2-5 µg/mL, Human
Immunocytochemistry/Immunofluorescence, 5 µg/mL, Human
Flow Cytometry (Fixed), 1-3 µg/1×10⁶ cells, Human
1. Dick, T., Ray, K., Salz, H. K., Chia, W. Cytoplasmic dynein (ddlc1) mutations cause morphogenetic defects and apoptotic cell death in Drosophila melanogaster. Molec. Cell. Biol.

Application Details

16: 1966-1977, 1996. 2. Fuhrmann, J. C., Kins, S., Rostaing, P., El Far, O., Kirsch, J., Sheng, M., Triller, A., Betz, H., Kneussel, M. Gephyrin interacts with dynein light chains 1 and 2, components of motor protein complexes. J. Neurosci. 22: 5393-5402, 2002. 3. He, Y. J., Meghani, K., Caron, M.-C., Yang, C., Ronato, D. A., Bian, J., Sharma, A., Moore, J., Niraj, J., Detappe, A., Doench, J. G., Legube, G., Root, D. E., D'Andrea, A. D., Drane, P., De, S., Konstantinopoulos, P. A., Masson, J.-Y., Chowdhury, D. DYNLL1 binds to MRE11 to limit DNA end resection in BRCA1-deficient cells. Nature 563: 522-526, 2018.

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na2HPO4.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.