

Datasheet for ABIN7599800
anti-PCDHa8 antibody (AA 118-801)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	PCDHa8
Binding Specificity:	AA 118-801
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PCDHa8 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS)

Product Details

Purpose:	Anti-PCDHA8 Antibody Picoband®
Immunogen:	E.coli-derived human PCDHA8 recombinant protein (Position: H118-Q801).
Isotype:	IgG
Cross-Reactivity (Details):	No cross reactivity with other proteins.
Characteristics:	Anti-PCDHA8 Antibody Picoband® (ABIN7599800). Tested in ELISA, WB, Flow Cytometry applications. This antibody reacts with Human, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	PCDHa8
Alternative Name:	PCDHA8 (PCDHa8 Products)
Background:	<p>Synonyms: 70 kDa ribosomal protein S6 kinase 1 antibody, KS6B1_HUMAN antibody, p70 alpha antibody, P70 beta 1 antibody, p70 ribosomal S6 kinase alpha antibody, p70 ribosomal S6 kinase beta 1 antibody, p70 S6 kinase alpha antibody, P70 S6 Kinase antibody, p70 S6 kinase alpha 1 antibody, p70 S6 kinase alpha 2 antibody, p70 S6K antibody, p70 S6K-alpha antibody, p70 S6KA antibody, p70(S6K) alpha antibody, p70(S6K)-alpha antibody, p70-alpha antibody, p70-S6K 1 antibody, p70-S6K antibody, P70S6K antibody, P70S6K1 antibody, p70S6Kb antibody, PS6K antibody, Ribosomal protein S6 kinase 70 kDa polypeptide 1 antibody, Ribosomal protein S6 kinase beta 1 antibody, Ribosomal protein S6 kinase beta-1 antibody, Ribosomal protein S6 kinase I antibody, RPS6KB1 antibody, S6K antibody, S6K-beta-1 antibody, S6K1 antibody, Serine/threonine kinase 14 alpha antibody, Serine/threonine-protein kinase 14A antibody, STK14A antibody</p> <p>Tissue Specificity: Expressed in all tissues.</p> <p>Background: This gene is a member of the protocadherin alpha gene cluster, one of three related gene clusters tandemly linked on chromosome five that demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The alpha gene cluster is composed of 15 cadherin superfamily genes related to the mouse CNR genes and consists of 13 highly similar and 2 more distantly related coding sequences. The tandem array of 15 N-terminal exons, or variable exons, are followed by downstream C-terminal exons, or constant exons, which are shared by all genes in the cluster. The large, uninterrupted N-terminal exons each encode six cadherin ectodomains while the C-terminal exons encode the cytoplasmic domain. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins that most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been observed and additional variants have been suggested but their full-length nature has yet to be determined.</p>
Molecular Weight:	150 kDa
Gene ID:	56140
UniProt:	Q9Y5H6

Application Details

Application Notes:	Western blot, 0.25-0.5 µg/mL, Human, Rat
	Flow Cytometry (Fixed), 1-3 µg/1x10 ⁶ cells, Human

Application Details

	ELISA, 0.1-0.5 µg/mL, - 1. Noonan, J. P., Li, J., Nguyen, L., Caoile, C., Dickson, M., Grimwood, J., Schmutz, J., Feldman, M. W., Myers, R. M. Extensive linkage disequilibrium, a common 16.7-kilobase deletion, and evidence of balancing selection in the human protocadherin alpha cluster. Am. J. Hum. Genet. 72: 621-635, 2003. 2. Ribich, S., Tasic, B., Maniatis, T. Identification of long-range regulatory elements in the protocadherin-alpha gene cluster. Proc. Nat. Acad. Sci. 103: 19719-19724, 2006. 3. Wu, Q., Maniatis, T. A striking organization of a large family of human neural cadherin-like cell adhesion genes. Cell 97: 779-790, 1999.
--	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.