

Datasheet for ABIN7599968
anti-NUP54 antibody (AA 132-465)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	NUP54
Binding Specificity:	AA 132-465
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NUP54 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Purpose:	Anti-NUP54 Antibody Picoband®
Immunogen:	E.coli-derived human NUP54 recombinant protein (Position: D132-Q465).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-NUP54 Antibody Picoband® (ABIN7599968). Tested in ELISA, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	NUP54
Alternative Name:	NUP54 (NUP54 Products)
Background:	<p>Synonyms: RNA-binding protein Nova-2, Astrocytic NOVA1-like RNA-binding protein, Neuro-oncological ventral antigen 2, NOVA2, ANOVA, NOVA3</p> <p>Tissue Specificity: Brain. Expression restricted to astrocytes.</p> <p>Background: Nucleoporin 54 (Nup54) is a protein that in humans is encoded by the NUP54 gene. The nuclear envelope creates distinct nuclear and cytoplasmic compartments in eukaryotic cells. It consists of two concentric membranes perforated by nuclear pores, large protein complexes that form aqueous channels to regulate the flow of macromolecules between the nucleus and the cytoplasm. These complexes are composed of at least 100 different polypeptide subunits, many of which belong to the nucleoporin family. This gene encodes a member of the phe-gly (FG) repeat-containing nucleoporin subset. Multiple alternatively spliced transcript variants have been found for this gene.</p>
Molecular Weight:	56 kDa
Gene ID:	53371
UniProt:	Q7Z3B4
Pathways:	SARS-CoV-2 Protein Interactome

Application Details

Application Notes:	<p>Western blot, 0.25-0.5 µg/mL, Human</p> <p>ELISA, 0.1-0.5 µg/mL, -</p> <p>1. Chug, H., Trakhanov, S., Hulsmann, B. B., Pleiner, T., Gorlich, D. Crystal structure of the metazoan Nup62-Nup58-Nup54 nucleoporin complex. Science 350: 106-110, 2015. 2. Hu, T., Guan, T., Gerace, L. Molecular and functional characterization of the p62 complex, an assembly of nuclear pore complex glycoproteins. J. Cell Biol. 134: 589-601, 1996.</p>
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL

Handling

Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.