

Datasheet for ABIN7599983
anti-PCBP2 antibody (AA 136-276)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	PCBP2
Binding Specificity:	AA 136-276
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PCBP2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC), Flow Cytometry (FACS)

Product Details

Purpose:	Anti-hnRNP E2/PCBP2 Antibody Picoband®
Immunogen:	E.coli-derived human hnRNP E2/PCBP2 recombinant protein (Position: D136-K276).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-hnRNP E2/PCBP2 Antibody Picoband® (ABIN7599983). Tested in ELISA, Flow Cytometry, IF, ICC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	PCBP2
Alternative Name:	PCBP2 (PCBP2 Products)
Background:	Synonyms: Peptidyl-prolyl cis-trans isomerase D, PPIase D Tissue Specificity: Widely expressed. Background: Poly(rC)-binding protein 2 is a protein that in humans is encoded by the PCBP2 gene. The protein encoded by this gene appears to be multifunctional. Along with PCBP-1 and hnRNPK, it is one of the major cellular poly(rC)-binding proteins. The encoded protein contains three K-homologous (KH) domains which may be involved in RNA binding. Together with PCBP-1, this protein also functions as a translational coactivator of poliovirus RNA via a sequence-specific interaction with stem-loop IV of the IRES, promoting poliovirus RNA replication by binding to its 5'-terminal cloverleaf structure. It has also been implicated in translational control of the 15-lipoxygenase mRNA, human papillomavirus type 16 L2 mRNA, and hepatitis A virus RNA. The encoded protein is also suggested to play a part in formation of a sequence-specific alpha-globin mRNP complex which is associated with alpha-globin mRNA stability. This multiexon structural mRNA is thought to be retrotransposed to generate PCBP-1, an intronless gene with functions similar to that of PCBP2. This gene and PCBP-1 have paralogous genes (PCBP3 and PCBP4) which are thought to have arisen as a result of duplication events of entire genes. This gene also has two processed pseudogenes (PCBP2P1 and PCBP2P2). Multiple transcript variants encoding different isoforms have been found for this gene.
Molecular Weight:	40 kDa
Gene ID:	5094
UniProt:	Q15366

Application Details

Application Notes:	Western blot, 0.1-0.25 µg/mL, Human, Mouse, Rat Immunocytochemistry/Immunofluorescence, 5 µg/mL, Human Flow Cytometry (Fixed), 1-3 µg/1x10⁶ cells, Human ELISA, 0.1-0.5 µg/mL, - 1. Bejerano, G., Lowe, C. B., Ahituv, N., King, B., Siepel, A., Salama, S. R., Rubin, E. M., Kent, W. J., Haussler, D. A distal enhancer and an ultraconserved exon are derived from a novel retroposon. Nature 441: 87-90, 2006. 2. Chkheidze, A. N., Liebhaber, S. A. A novel set of nuclear localization signals determine distributions of the alpha-CP RNA-binding proteins. Molec. Cell. Biol. 23: 8405-8415, 2003. 3. Eiring, A. M., Harb, J. G., Neviani, P., Garton, C., Oaks, J. J., Spizzo, R., Liu, S.,
--------------------	--

Application Details

Schwind, S., Santhanam, R., Hickey, C. J., Becker, H., Chandler, J. C., and 13 others. miR-328 functions as an RNA decoy to modulate hnRNP E2 regulation of mRNA translation in leukemic blasts. Cell 140: 652-665, 2010.

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.