

Datasheet for ABIN7600094 anti-PRPF39 antibody (AA 149-664)



Overview

Quantity:	100 μg
Target:	PRPF39
Binding Specificity:	AA 149-664
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRPF39 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC), Flow Cytometry (FACS)

Product Details

Purpose:	Anti-PRPF39 Antibody Picoband®
Immunogen:	E.coli-derived human PRPF39 recombinant protein (Position: D149-Y664).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-PRPF39 Antibody Picoband® (ABIN7600094). Tested in ELISA, Flow Cytometry, IF, ICC,
	WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates
	this is a premium antibody that guarantees superior quality, high affinity, and strong signals
	with minimal background in Western blot applications. Only our best-performing antibodies are
	designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	PRPF39
Alternative Name:	PRPF39 (PRPF39 Products)
Background:	Synonyms: Kelch repeat and BTB domain-containing protein 2, BTB and kelch domain-containing protein 1, KBTBD2, BKLHD1, KIAA1489 Tissue Specificity: Detected in liver, skeletal muscle, kidney, pancreas, spleen, thyroid, testis, ovary, small intestine and colon. Background: Predicted to be involved in mRNA 5'-splice site recognition. Predicted to be part of U1 snRNP, U2-type prespliceosome, and commitment complex.
Molecular Weight:	90 kDa
Gene ID:	55015

Application Details

Application Notes:	Western blot, 0.25-0.5 μg/mL, Human, Mouse, Rat
	Immunocytochemistry/Immunofluorescence, 5 µg/mL, Human
	Flow Cytometry (Fixed), 1-3 µg/1x10 ⁶ cells, Human
	ELISA, 0.1-0.5 μg/mL, -
	1. Hartz, P. A. Personal Communication. Baltimore, Md. 11/2/2012. 2. Stark, A. L., Delaney, S.
	M., Wheeler, H. E., Im, H. K., Dolan, M. E. Functional consequences of PRPF39 on distant genes
	and cisplatin sensitivity. Hum. Molec. Genet. 21: 4348-4355, 2012.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 μg/mL.
Concentration:	500 μg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.