

# Datasheet for ABIN7600367 anti-SMARCA2 antibody (AA 181-624)



#### Go to Product page

_				
( )	1//	rv	IO	Λ/
( )	VC	. I V	1	v v

Quantity:	100 μg
Target:	SMARCA2
Binding Specificity:	AA 181-624
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SMARCA2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS)

### **Product Details**

Purpose:	Anti-SMARCA2/BRM Antibody Picoband®
Immunogen:	E.coli-derived human SMARCA2/BRM recombinant protein (Position: Q181-N624).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-SMARCA2/BRM Antibody Picoband® (ABIN7600367). Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

## **Target Details**

Target:	SMARCA2	
Alternative Name:	SMARCA2 (SMARCA2 Products)	
Background:	Synonyms: Probable global transcription activator SNF2L2, ATP-dependent helicase SMARCA2	
	BRG1-associated factor 190B, BAF190B, Protein brahma homolog, hBRM, SNF2-alpha,	
	SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily A	
	member 2, SMARCA2, BAF190B, BRM, SNF2A, SNF2L2	
	Tissue Specificity: Detected in milk (at protein level)	
	Background: Probable global transcription activator SNF2L2 is a protein that in humans is	
	encoded by the SMARCA2 gene. It is mapped to 9p24.3. The protein encoded by this gene is a	
	member of the SWI/SNF family of proteins and is highly similar to the brahma protein of	
	Drosophila. Members of this family have helicase and ATPase activities and are thought to	
	regulate transcription of certain genes by altering the chromatin structure around those genes.	
	The encoded protein is part of the large ATP-dependent chromatin remodeling complex	
	SNF/SWI, which is required for transcriptional activation of genes normally repressed by	
	chromatin. Alternatively spliced transcript variants encoding different isoforms have been	
	found for this gene, which contains a trinucleotide repeat (CAG) length polymorphism.	
Molecular Weight:	210 kDa	
Gene ID:	6595	
UniProt:	P51531	
Application Details		
Application Notes:	Western blot, 0.25-0.5 μg/mL, Human, Mouse, Rat	
	Flow Cytometry (Fixed), 1-3 µg/1x10 <sup>6</sup> cells, Human	
	ELISA, 0.1-0.5 μg/mL, -	
	1. de la Serna, I. L., Carlson, K. A., Imbalzano, A. N. Mammalian SWI/SNF complexes promote	
	MyoD-mediated muscle differentiation. Nature Genet. 27: 187-190, 2001. 2. Hakimi, MA.,	
	Bochar, D. A., Schmiesing, J. A., Dong, Y., Barak, O. G., Speicher, D. W., Yokomori, K., Shiekhattar	
	R. A chromatin remodelling complex that loads cohesin onto human chromosomes. Nature	
	418: 994-998, 2002.	
Restrictions:	For Research Use only	
Handling		

## Handling

Reconstitution:	Add 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 μg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4, 0.05 mg Sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.