

Datasheet for ABIN7600639  
**anti-PPP1R15B antibody (AA 211-657)**



[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	PPP1R15B
Binding Specificity:	AA 211-657
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PPP1R15B antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS)

## Product Details

Purpose:	Anti-PPP1R15B Antibody Picoband®
Immunogen:	E.coli-derived human PPP1R15B recombinant protein (Position: Q211-D657).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins
Characteristics:	Anti-PPP1R15B Antibody Picoband® (ABIN7600639). Tested in ELISA, WB, Flow Cytometry applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

## Target Details

Target:	PPP1R15B
Alternative Name:	PPP1R15B ( <a href="#">PPP1R15B Products</a> )
Background:	<p>Synonyms: Protein NDRG3,N-myc downstream-regulated gene 3 protein,NDRG3,</p> <p>Tissue Specificity: Ubiquitous. Highly expressed in brain. .</p> <p>Background: PPP1R15B(Protein phosphatase 1, regulatory subunit 15b), also called CREP, promotes dephosphorylation of the transcription initiation factor EIF2-alpha through recruitment of protein phosphatase-1(PP1) catalytic subunits. The PPP1R15B gene is mapped to chromosome 1q32.1 based on an alignment of thePPP1R15B sequence by Hartz(2010). Harding et al.(2009) obtained Ppp1r15b -/- mice at a mendelian ratio. However, Ppp1r15b -/- newborns were half the size of their wildtype littermates, were notably pale, and failed to nurse, and none survived the first day of postnatal life. Ppp1r15b -/- embryos that were also homozygous for an Eif2-alpha mutation that prevented Eif2-alpha phosphorylation were normalized, including elevated birth size and restored red blood cell count, compared with Ppp1r15b -/- embryos with wildtype Eif2-alpha.</p>
Molecular Weight:	110 kDa
Gene ID:	84919
Pathways:	<a href="#">ER-Nucleus Signaling</a>

## Application Details

Application Notes:	<p>Western blot, 0.25-0.5 µg/mL, Human</p> <p>Flow Cytometry (Fixed), 1-3 µg/1x10<sup>6</sup> cells, Human</p> <p>ELISA, 0.1-0.5 µg/mL, -</p> <p>1. Harding, H. P., Zhang, Y., Scheuner, D., Chen, J.-J., Kaufman, R. J., Ron, D. Ppp1r15 gene knockout reveals an essential role for translation initiation factor 2 alpha (eIF2-alpha) dephosphorylation in mammalian development. Proc. Nat. Acad. Sci. 106: 1832-1837, 2009.</p>
Restrictions:	For Research Use only

## Handling

Format:	Lyophilized
Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.

## Handling

---

Storage: 4 °C, -20 °C

---

Storage Comment: At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month.  
It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.