

Datasheet for ABIN7600892  
**anti-PDE6B antibody (AA 25-237)**



[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	PDE6B
Binding Specificity:	AA 25-237
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This PDE6B antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC), Flow Cytometry (FACS)

## Product Details

Purpose:	Anti-PDE6 beta/PDE6B Antibody Picoband® (monoclonal, 8I2D7)
Immunogen:	E.coli-derived human PDE6 beta/PDE6B recombinant protein (Position: K25-Q237).
Clone:	8I2D7
Isotype:	IgG2a
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-PDE6 beta/PDE6B Antibody Picoband® (monoclonal, 8I2D7) (ABIN7600892). Tested in Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.

## Product Details

Purification: Immunogen affinity purified.

## Target Details

Target: PDE6B

Alternative Name: PDE6B ([PDE6B Products](#))

Background: Synonyms: Arylsulfatase A, ASA, Cerebroside-sulfatase, Arylsulfatase A component B, Arylsulfatase A component C, ARSA  
Background: Photon absorption triggers a signaling cascade in rod photoreceptors that activates cGMP phosphodiesterase (PDE), resulting in the rapid hydrolysis of cGMP, closure of cGMP-gated cation channels, and hyperpolarization of the cell. PDE is a peripheral membrane heterotrimeric enzyme made up of alpha, beta, and gamma subunits. This gene encodes the beta subunit. Mutations in this gene result in retinitis pigmentosa and autosomal dominant congenital stationary night blindness. Multiple transcript variants encoding different isoforms have been found for this gene.

Molecular Weight: 98 kDa

Gene ID: 5158

UniProt: [P35913](#)

Pathways: [Regulation of G-Protein Coupled Receptor Protein Signaling, Phototransduction](#)

## Application Details

Application Notes: Western blot, 0.25-0.5 µg/mL, Mouse, Rat  
Immunohistochemistry(Paraffin-embedded Section), 2-5 µg/mL, Rat  
Immunocytochemistry/Immunofluorescence, 5 µg/mL, Human  
Flow Cytometry (Fixed), 1-3 µg/1x<sup>6</sup> cells, Human  
1. Hmani-Aifa, M., Benzina, Z., Zulficar, F., Dhouib, H., Shahzadi, A., Ghorbel, A., Rebai, A., Soderkvist, P., Riazuddin, S., Kimberling, W. J., Ayadi, H. Identification of two new mutations in the GPR98 and the PDE6B genes segregating in a Tunisian family. *Europ. J. Hum. Genet.* 17: 474-482, 2009. 2. Tsang, S. H., Woodruff, M. L., Jun, L., Mahajan, V., Yamashita, C. K., Pedersen, R., Lin, C.-S., Goff, S. P., Rosenberg, T., Larsen, M., Farber, D. B., Nusinowitz, S. Transgenic mice carrying the H258N mutation in the gene encoding the beta-subunit of phosphodiesterase-6 (PDE6B) provide a model for human congenital stationary night blindness. *Hum. Mutat.* 28: 243-254, 2007. 3. Zhao, L., Zabel, M. K., Wang, X., Ma, W., Shah, P., Fariss, R. N., Qian, H., Parkhurst, C. N., Gan, W.-B., Wong, W. T. Microglial phagocytosis of living photoreceptors contributes to

## Application Details

---

inherited retinal degeneration. EMBO Molec. Med. 7: 1179-1197, 2015.

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

Reconstitution: Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.

Concentration: 500 µg/mL

Buffer: Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na<sub>2</sub>HPO<sub>4</sub>.

Storage: 4 °C, -20 °C

Storage Comment: At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month.  
It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.