

Datasheet for ABIN7601094  
**anti-MUS81 antibody (AA 283-483)**



[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	MUS81
Binding Specificity:	AA 283-483
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MUS81 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB)

## Product Details

Purpose:	Anti-MUS81 Antibody Picoband®
Immunogen:	E.coli-derived human MUS81 recombinant protein (Position: H283-R483).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-MUS81 Antibody Picoband® (ABIN7601094). Tested in ELISA, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

## Target Details

Target:	MUS81
Alternative Name:	MUS81 ( <a href="#">MUS81 Products</a> )
Background:	<p>Synonyms: Lipopolysaccharide-binding protein, LBP, Lbp,</p> <p>Tissue Specificity: Detected in blood serum (at protein level).</p> <p>Background: Crossover junction endonuclease MUS81 is an enzyme that in humans is encoded by the MUS81 gene. This gene encodes a structure-specific endonuclease which belongs to the XPF/MUS81 endonuclease family and plays a critical role in the resolution of recombination intermediates during DNA repair after inter-strand cross-links, replication fork collapse, and DNA double-strand breaks. The encoded protein associates with one of two closely related essential meiotic endonuclease proteins (EME1 or EME2) to form a complex that processes DNA secondary structures. It contains an N-terminal DEAH helicase domain, an excision repair cross complementation group 4 (ERCC4) endonuclease domain, and two tandem C-terminal helix-hairpin-helix domains. Mice with a homozygous knockout of the orthologous gene have significant meiotic defects including the failure to repair a subset of DNA double strand breaks.</p>
Molecular Weight:	52-72 kDa
Gene ID:	80198
Pathways:	<a href="#">DNA Damage Repair</a>

## Application Details

Application Notes:	<p>Western blot, 0.25-0.5 µg/mL, Human, Mouse, Rat</p> <p>ELISA, 0.1-0.5 µg/mL, -</p> <p>1. Chen, X.-B., Melchionna, R., Denis, C.-M., Gaillard, P.-H. L., Blasina, A., Van de Weyer, I., Boddy, M. N., Russell, P., Vialard, J., McGowan, C. H. Human Mus81-associated endonuclease cleaves Holliday junctions in vitro. <i>Molec. Cell</i> 8: 1117-1127, 2001. 2. Ciccia, A., Constantinou, A., West, S. C. Identification and characterization of the human Mus81-Eme1 endonuclease. <i>J. Biol. Chem.</i> 278: 25172-25178, 2003. 3. Ciccia, A., Ling, C., Coulthard, R., Yan, Z., Xue, Y., Meetei, A. R., Laghmani, E. H., Joenje, H., McDonald, N., de Winter, J. P., Wang, W., West, S. C. Identification of FAAP24, a Fanconi anemia core complex protein that interacts with FANCM. <i>Molec. Cell</i> 25: 331-343, 2007.</p>
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized

## Handling

---

Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage:	4 °C, -20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.